

Quantum Near Horizon Geometry of Black 0-Brane

Yoshifumi Hyakutake

*College of Science, Ibaraki University
Bunkyo 1-1, Mito, Ibaraki 310-0062, Japan*

Abstract

We investigate a bunch of D0-branes to reveal its quantum nature from the gravity side. In the classical limit, it is well described by a non-extremal black 0-brane in type IIA supergravity. The solution is uplifted to the eleven dimensions and expressed by a non-extremal M-wave solution. After reviewing the effective action for the M-theory, we explicitly solve the equations of motion for the near horizon geometry of the M-wave. As a result we derive a unique solution which includes the effect of the quantum gravity. Thermodynamic property of the quantum near horizon geometry of the black 0-brane is also studied by using Wald's formula. Combining our result with that of the Monte Carlo simulation of the dual thermal gauge theory, we find strong evidence for the gauge/gravity duality in the D0-branes system at the level of quantum gravity.

1 Introduction

Superstring theory is a promising candidate for the theory of quantum gravity, and it plays important roles to reveal quantum nature of black holes. Fundamental objects in the superstring theory are D-branes as well as strings [1], and in the low energy limit their dynamics are governed by supergravity. The D-branes are described by classical solutions in the supergravity, which are called black branes [2, 3]. A special class of them has event horizon like the black holes and its entropy can be evaluated by the area law. Interestingly the entropy can be statistically explained by counting number of microstates in the gauge theory on the D-branes [4]. This motivates us to study the black hole thermodynamics from the gauge theory. Furthermore it is conjectured that the near horizon geometry of the black brane corresponds to the gauge theory on the D-branes [5]. If this gauge/gravity duality is correct, the strong coupling limit of the gauge theory can be analyzed by the supergravity [6, 7].

In this paper we consider a bunch of D0-branes in type IIA superstring theory. In the low energy limit, a bunch of D0-branes with additional internal energy are well described by non-extremal black 0-brane solution in type IIA supergravity [2, 3]. After taking near horizon limit, the metric becomes AdS black hole like geometry in ten dimensional space-time [8]. From the gauge/gravity duality, this geometry corresponds to the strong coupling limit of the gauge theory on the D0-branes [8], which is described by (1+0)-dimensional $U(N)$ super Yang-Mills theory [9]. This gauge theory is paid much attention as nonperturbative definition of M-theory [10, 11], which is the strong coupling description of the type IIA superstring theory [12, 13]. Recently nonperturbative aspects of the gauge theory are studied by the computer simulation [14]-[24]. (See refs. [25], [26] for reviews including other topics.) Especially in ref. [19], physical quantities of the thermal gauge theory, such as the internal energy, are evaluated numerically, and a direct test of the gauge/gravity duality is performed including α' correction to the type IIA supergravity. Furthermore, if the internal energy of the black 0-brane can be evaluated precisely from the gravity side including g_s correction, it is possible to give a direct test for the gauge/gravity duality at the level of quantum gravity [24]. ($\alpha' = \ell_s^2$ is the string length squared and g_s is the string coupling constant.)

The purpose of this paper is to derive quantum correction to the near horizon geometry of the non-extremal black 0-brane directly from the gravity side. In order to do this, we need to know an effective action which include quantum correction to the type IIA supergravity. In principle the effective action can be constructed so as to be consistent with the scattering amplitudes in the type IIA superstring theory [27], and it is expressed by double expansion of α' and g_s . For example, since four point amplitudes of gravitons at tree and one loop level are nontrivial, there should exist terms like $\alpha'^3 e^{-2\phi} t_8 t_8 R^4$ and $\alpha'^3 g_s^2 t_8 t_8 R^4$ in the effective

action, respectively [27]–[36]. These are called higher derivative terms and t_8 represents products of four Kronecker’s deltas with eight indices. Especially we are interested in the latter terms, which give nontrivial g_s corrections to the geometry. These higher derivative terms often play important roles to count the entropy of extremal black holes [37, 38].

It is necessary that the effective action of the type IIA superstring should possess local supersymmetry in ten dimensions. So the supersymmetrization of $\alpha'^3 g_s^2 t_8 R^4$ is very important [28, 29, 30, 33, 35, 36] to understand the structure of effective action. Although the task is not completed yet, since our interest is on the geometry of the black 0-brane, it is enough to know terms which contain the metric, dilaton field and R-R 1-form field only. Notice that these fields are collected into the metric in eleven dimensional supergravity [39], and the black 0-brane is expressed by M-wave solution. Then $\alpha'^3 g_s^2 t_8 R^4$ and other terms which include the dilaton and R-R 1-form field are simply collected into $\ell_p^6 t_8 R^4$ terms in eleven dimensions. Here $\ell_p = \ell_s g_s^{1/3}$ is the Planck length in eleven dimensions. Thus we consider the effective action for the M-theory and investigate quantum corrections to the near horizon geometry of the non-extremal M-wave. We show equations of motion for the effective action and explicitly solve them up to the order of g_s^2 . The M-wave geometry receives the quantum corrections and thermodynamic quantities for the M-wave are modified. Especially the internal energy of the M-wave is obtained quantitatively including quantum effect of the gravity.

Organization of the paper is as follows. In section 2, we review the classical near horizon geometry of the black 0-brane in ten dimensions, and uplift it to that of the M-wave in eleven dimensions. In section 3, we discuss the higher derivative corrections in the type IIA superstring theory and the M-theory, and solve the equations of motion for the near horizon geometry of the non-extremal M-wave in section 4. In section 5, we evaluate the entropy and the energy of the M-wave up to $1/N^2$. Section 6 is devoted to conclusion and discussion. Detailed calculations and discussions on the ambiguities of the higher derivative corrections are collected in the appendices.

2 Classical Near Horizon Geometry of Black 0-Brane

In this section, we briefly review the non-extremal solution of the black 0-brane which carries mass and R-R charge. Especially we up lift the solution to eleven dimensions and show that the black 0-brane is described by the M-wave solution.

In the low energy limit, the dynamics of massless modes in type IIA superstring theory are governed by type IIA supergravity. Since we are interested in the black 0-brane which couples to the graviton $g_{\mu\nu}$, the dilaton ϕ and R-R 1-form field C_μ , the relevant part of the

type IIA supergravity action is given by

$$S_{10}^{(0)} = \frac{1}{2\kappa_{10}^2} \int d^{10}x \sqrt{-g} \left\{ e^{-2\phi} (R + 4\partial_\mu \phi \partial^\mu \phi) - \frac{1}{4} G_{\mu\nu} G^{\mu\nu} \right\}, \quad (1)$$

where $2\kappa_{10}^2 = (2\pi)^7 \ell_s^8 g_s^2$ and $G_{\mu\nu}$ is the field strength of C_μ . g_s and ℓ_s are the string coupling constant and the string length, respectively. It is possible to solve the equations of motion by making the ansatz that the metric is static and has SO(9) rotation symmetry. Then we obtain non-extremal solution of the black 0-brane. (See ref. [40] for example.)

$$\begin{aligned} ds_{10}^2 &= -\tilde{H}^{-\frac{1}{2}} \tilde{F} dt^2 + \tilde{H}^{\frac{1}{2}} \tilde{F}^{-1} dr^2 + \tilde{H}^{\frac{1}{2}} r^2 d\Omega_8^2, \\ e^\phi &= \tilde{H}^{\frac{3}{4}}, \quad C = \left(\frac{r_+}{r_-} \right)^{\frac{7}{2}} \tilde{H}^{-1} dt, \\ \tilde{H} &= 1 + \frac{r_-^7}{r^7}, \quad \tilde{F} = 1 - \frac{r_+^7 - r_-^7}{r^7}. \end{aligned} \quad (2)$$

The horizon is located at $r_H = (r_+^7 - r_-^7)^{\frac{1}{7}}$. Parameters r_\pm are related to the mass M_0 and the R-R charge Q_0 of the black 0-brane by

$$M_0 = \frac{V_{S^8}}{2\kappa_{10}^2} (8r_+^7 - r_-^7), \quad Q_0 = \frac{N}{\ell_s g_s} = \frac{7V_{S^8}}{2\kappa_{10}^2} (r_+ r_-)^{\frac{7}{2}}, \quad (3)$$

where N is a number of D0-branes and $V_{S^8} = \frac{2\pi^{9/2}}{\Gamma(9/2)} = \frac{2(2\pi)^4}{7 \cdot 15}$ is the volume of S^8 . Now the parameters r_\pm are expressed as

$$r_\pm^7 = (1 + \delta)^{\pm 1} (2\pi)^2 15\pi g_s N \ell_s^7, \quad (4)$$

where δ is a non-negative parameter. The extremal limit $r_+ = r_-$ is saturated when $\delta = 0$.

Let us rewrite the solution (2) in terms of $U = r/\ell_s^2$ and $\lambda = g_s N / (2\pi)^2 \ell_s^3$, which correspond to typical energy scale and 't Hooft coupling in the dual gauge theory, respectively. The near horizon limit of the non-extremal black 0-brane is taken by $\ell_s \rightarrow 0$ while U , λ and δ/ℓ_s^4 are fixed. Then the near horizon limit of the solution (2) becomes [8]

$$\begin{aligned} ds_{10}^2 &= \ell_s^2 \left(-H^{-\frac{1}{2}} F dt^2 + H^{\frac{1}{2}} F^{-1} dU^2 + H^{\frac{1}{2}} U^2 d\Omega_8^2 \right), \\ e^\phi &= \ell_s^{-3} H^{\frac{3}{4}}, \quad C = \ell_s^4 H^{-1} dt, \\ H &= \frac{(2\pi)^4 15\pi \lambda}{U^7}, \quad F = 1 - \frac{U_0^7}{U^7}, \end{aligned} \quad (5)$$

where $U_0^7 = \frac{2\delta}{\ell_s^4} (2\pi)^4 15\pi \lambda$.

The type IIA supergravity is related to the eleven dimensional supergravity via circle compactification. In fact, the eleven dimensional metric is related to the ten dimensional one like $ds_{11}^2 = e^{-2\phi/3} ds_{10}^2 + e^{4\phi/3} (dz - C_\mu dx^\mu)^2$. The near horizon limit of the non-extremal solution of the black 0-brane (5) can be up lifted to eleven dimensions as

$$ds_{11}^2 = \ell_s^4 \left(-H^{-1} F dt^2 + F^{-1} dU^2 + U^2 d\Omega_8^2 + (\ell_s^{-4} H^{\frac{1}{2}} dz - H^{-\frac{1}{2}} dt)^2 \right). \quad (6)$$

This represents the near horizon limit of the non-extremal M-wave solution in eleven dimensions. The solution is purely geometrical and the expressions become simple. Furthermore, on the geometrical part, quantum corrections to the eleven dimensional supergravity are under control. This is the reason why we execute analyses of the solution in eleven dimensions.

3 Quantum Correction to Eleven Dimensional Supergravity

The eleven dimensional supergravity is realized as the low energy limit of the M-theory. A fundamental object in the M-theory is a membrane and if we could take account of interaction of membranes, the effective action of the M-theory would become the eleven dimensional supergravity with some higher derivative terms. Unfortunately quantization of the membrane has not been completed so far. It is, however, possible to derive the relevant part of the quantum corrections in the M-theory by requiring local supersymmetry. In this section we review the quantum corrections to the eleven dimensional supergravity.

Massless fields of the eleven dimensional supergravity consists of a vielbein $e^a{}_\mu$, a Majorana gravitino ψ_μ and a 3-form field $A_{\mu\nu\rho}$. Since we are only interested in the M-wave solution, we only need to take account of the action which only depends on the graviton.

$$2\kappa_{11}^2 S_{11}^{(0)} = \int d^{11}x eR, \quad (7)$$

where $2\kappa_{11}^2 = (2\pi)^8 \ell_p^9 = (2\pi)^8 \ell_s^9 g_s^3$. Notice that after the dimensional reduction this becomes the action (1), which contains the dilation and the R-R 1-form field as well as the graviton in ten dimensions [39].

Of course there are other terms which depend on ψ_μ and $A_{\mu\nu\rho}$, which are completely determined by the local supersymmetry. For example, a variation of the vielbein under the local supersymmetry is given by $\delta[e] = [\bar{\epsilon}\psi]$. Here we use a symbol $[X]$ to abbreviate indices and gamma matrices in X , and ϵ represents a parameter of the local supersymmetry. Then the variation of the scalar curvature is written by $\delta[eR] = [eR\bar{\epsilon}\psi]$. In order to cancel this, we see that a variation of the Majorana gravitino should include $\delta[\psi] = [D\epsilon] + \dots$ and simultaneously there should exist a term like $[e\bar{\psi}\psi_2]$ in the action. Here ψ_2 represents the field strength of the Majorana gravitino. By continuing this process, it is possible to determine the structure of the 11 dimensional supergravity completely [39].

Now let us discuss quantum corrections to the eleven dimensional supergravity. Since the M-theory is related to the type IIA superstring theory by the dimensional reduction, the effective action of the M-theory should contain that of the type IIA superstring theory. The latter can be obtained so as to be consistent with scattering amplitudes of strings, and it is well-known that leading corrections to the type IIA supergravity include terms

like $[eR^4]$. This is directly uplifted to the eleven dimensions and we see that the effective action of the M-theory should include terms like $B_1 = [eR^4]_7$. The subscript 7 indicates that there are potentially 7 independent terms if we consider possible contractions of 16 indices out of 4 Riemann tensors. (To be more precise, we excluded terms which contain Ricci tensor or scalar curvature, since these can be eliminated by redefinition of the graviton. Discussions on these terms will be found in the appendix C.) As in the case of the eleven dimensional supergravity, it is possible to determine other corrections by requiring the local supersymmetry. For example, variations of B_1 under the local supersymmetry contain terms like $V_1 = [eR^4\bar{\epsilon}\psi]$. In order to cancel these terms, $B_{11} = [e\epsilon_{11}AR^4]_2$ and $F_1 = [eR^3\bar{\psi}\psi_2]_{92}$ should exist in the action. The structures of B_1 , B_{11} and F_1 are severely restricted by the local supersymmetry. By continuing this process, it is possible to show that a combination of terms in B_1 are completely determined up to over all factor [35, 36]. The result become as follows.

$$\begin{aligned}
2\kappa_{11}^2 S_{11}^{(1)} &= \frac{\pi^2 \ell_p^6}{3 \cdot 2^8 4!} \int d^{11}x e \left(t_8 t_8 R^4 - \frac{1}{4!} \epsilon_{11} \epsilon_{11} R^4 \right) \\
&= \frac{\pi^2 \ell_p^6}{3 \cdot 2^8 4!} \int d^{11}x e \left\{ 24 (R_{abcd} R_{abcd} R_{efgh} R_{efgh} - 64 R_{abcd} R_{aefg} R_{bcdh} R_{efgh} \right. \\
&\quad + 2 R_{abcd} R_{abef} R_{cdgh} R_{efgh} + 16 R_{abcd} R_{aebf} R_{cgdh} R_{efgh} \\
&\quad \left. - 16 R_{abcd} R_{aefg} R_{befh} R_{cdgh} - 16 R_{abcd} R_{aefg} R_{bfhe} R_{cdgh} \right\}. \quad (8)
\end{aligned}$$

Here t_8 is products of four Kronecker's deltas with eight indices and ϵ_{11} is an antisymmetric tensor with eleven indices. Local Lorentz indices are labelled by $a, b, \dots = 0, 1, \dots, 10$. Although all indices are lowered, it is understood those are contracted by the flat metric η_{ab} . The Riemann tensor with local Lorentz indices is defined by $R_{abcd} = e^\mu{}_c e^\nu{}_d (\partial_\mu \omega_{\nu ab} - \partial_\nu \omega_{\mu ab} + \omega_{\mu a}{}^\epsilon \omega_{\nu b \epsilon} - \omega_{\nu a}{}^\epsilon \omega_{\mu b \epsilon})$, where $\omega_{\mu ab}$ is a spin connection and μ, ν are space-time indices. The over all factor in eq. (8) is determined by employing the result of 1-loop four graviton amplitude in the type IIA superstring theory.

Since the near horizon limit of the M-wave solution (6) is purely geometrical, it is possible to examine the leading quantum corrections to it from the action (8). Other terms which depend on the 3-form field are irrelevant to the analyses for the M-wave. In summary the effective action of the M-theory is described by

$$S_{11} = S_{11}^{(0)} + S_{11}^{(1)} = \frac{1}{2\kappa_{11}^2} \int d^{11}x e \left\{ R + \gamma \ell_s^{12} \left(t_8 t_8 R^4 - \frac{1}{4!} \epsilon_{11} \epsilon_{11} R^4 \right) \right\}, \quad (9)$$

where $\gamma = \frac{\pi^2}{3 \cdot 2^8 4!} \frac{g_s^2}{\ell_s^6} = \frac{\pi^6}{2^7 3^2} \frac{\lambda^2}{N^2}$. Notice that the parameter γ remains finite after the decoupling limit is taken. After the dimensional reduction, the action (9) becomes the effective action of the type IIA superstring theory, which includes the 1-loop effect of the gravity.

Now we derive equations of motion for the action (9). Although the derivation is straightforward, we need to labor at many calculations because of the higher derivative terms in the action. Therefore in practice we use the Mathematica code for the calculations. Below we show the points of the calculations to build the code.

First of all we list variations of the fields with respect to the vielbein.

$$\begin{aligned}
\delta e &= -e e^i{}_\mu \delta e^\mu{}_i = -e \eta_{ij} \delta e^{ij}, \\
\delta \omega_{cab} &= e^\rho{}_c \delta \omega_{\rho ab} = (\delta^k_{[a} \eta_{b]i} \eta_{cj} + \delta^k_{[a} \eta_{b]j} \eta_{ci} + \delta^k_c \eta_{i[a} \eta_{b]j}) D_k \delta e^{ij}, \\
\delta R_{abcd} &= \delta e^\mu{}_c R_{ab\mu d} + \delta e^\mu{}_d R_{abc\mu} + e^\mu{}_c e^\nu{}_d \delta R_{ab\mu\nu} = -2\delta e^{ij} R_{abi[c} \eta_{d]j} + 2D_{[c} \delta \omega_{d]ab}, \\
\delta R_{ab} &= -\delta e^{ij} R_{ajib} + \delta e^{ij} R_{ai} \eta_{bj} + D_b \delta \omega^c{}_{ac} - D_c \delta \omega_{ba}{}^c,
\end{aligned} \tag{10}$$

where $\delta e^{ij} \equiv e^i{}_\mu \delta e^{\mu j}$. Then variations of the higher derivative terms are evaluated as

$$\begin{aligned}
& e \delta \left(t_8 t_8 R^4 - \frac{1}{4!} \epsilon_{11} \epsilon_{11} R^4 \right) \\
&= 24e \left\{ 4(\delta R_{abcd}) R_{abcd} R_{efgh} R_{efgh} - 64(\delta R_{abcd}) R_{abce} R_{dfgh} R_{efgh} \right. \\
&\quad + 8(\delta R_{abcd}) R_{abef} R_{cdgh} R_{efgh} + 64(\delta R_{abcd}) R_{aecg} R_{bf dh} R_{efgh} \\
&\quad - 64(\delta R_{abcd}) R_{abeg} R_{cf eh} R_{dfgh} - 64(\delta R_{abcd}) R_{efag} R_{efch} R_{gbhd} \\
&\quad \left. + 32(\delta R_{abcd}) R_{abef} R_{cegh} R_{dfgh} \right\} \\
&= e(\delta R_{abcd}) X_{abcd} \\
&= 2e \delta e^{ij} R_{abci} X^{abc}{}_j - 2e X^{abcd} D_d \delta \omega_{cab} \\
&\cong 2e \delta e^{ij} R_{abci} X^{abc}{}_j - 2e (\delta^k_a \eta_{bi} \eta_{cj} + \delta^k_a \eta_{bj} \eta_{ci} + \delta^k_c \eta_{ia} \eta_{bj}) D_k D_d X^{abcd} \delta e^{ij} \\
&= 2e R_{abci} X^{abc}{}_j \delta e^{ij} - 2e D^c D^d (X_{cijd} + X_{cjid} + X_{ijcd}) \delta e^{ij} \\
&= e(3R_{abci} X^{abc}{}_j - R_{abcj} X^{abc}{}_i) \delta e^{ij} - 2e D^c D^d (X_{cijd} + X_{cjid}) \delta e^{ij} \\
&= e(3R_{abci} X^{abc}{}_j - R_{abcj} X^{abc}{}_i - 4D_{(a} D_b) X^a{}_{ij}{}^b) \delta e^{ij},
\end{aligned} \tag{11}$$

where we defined

$$\begin{aligned}
X_{abcd} &= \frac{1}{2} (X'_{[ab][cd]} + X'_{[cd][ab]}), \\
X'_{abcd} &= 96 (R_{abcd} R_{efgh} R_{efgh} - 16 R_{abce} R_{dfgh} R_{efgh} + 2 R_{abef} R_{cdgh} R_{efgh} \\
&\quad + 16 R_{aecg} R_{bf dh} R_{efgh} - 16 R_{abeg} R_{cf eh} R_{dfgh} - 16 R_{efag} R_{efch} R_{gbhd} \\
&\quad + 8 R_{abef} R_{cegh} R_{dfgh}).
\end{aligned} \tag{12}$$

Finally we obtain the equations of motion for the effective action (9).

$$\begin{aligned}
E_{ij} &\equiv R_{ij} - \frac{1}{2} \eta_{ij} R + \gamma \ell_s^{12} \left\{ -\frac{1}{2} \eta_{ij} \left(t_8 t_8 R^4 - \frac{1}{4!} \epsilon_{11} \epsilon_{11} R^4 \right) \right. \\
&\quad \left. + \frac{3}{2} R_{abci} X^{abc}{}_j - \frac{1}{2} R_{abcj} X^{abc}{}_i - 2D_{(a} D_b) X^a{}_{ij}{}^b \right\} = 0.
\end{aligned} \tag{13}$$

As mentioned before the action (9) is not unique due to the ambiguity of field redefinitions, such as $g_{\mu\nu} \rightarrow g'_{\mu\nu} = g_{\mu\nu} + \gamma \ell_s^{12} R^2 R_{\mu\nu}$. Therefore the equations of motion are not unique as well. We will discuss, however, that physical quantities of the M-wave do not depend on these ambiguities. (See appendix D.)

4 Quantum Near Horizon Geometry of Black 0-Brane

In the previous section, we have explained the effective action of the M-theory (9), and derived the equations of motion (13). In this section we solve them up to the linear order of γ and obtain the non-extremal solution of the M-wave with quantum gravity correction.

In order to obtain the solution of (13), we relax the ansatz for the M-wave as

$$ds_{11}^2 = \ell_s^4 \left(-H_1^{-1} F_1 dt^2 + F_1^{-1} U_0^2 dx^2 + U_0^2 x^2 d\Omega_8^2 + (\ell_s^{-4} H_2^{\frac{1}{2}} dz - H_3^{-\frac{1}{2}} dt)^2 \right), \quad (14)$$

$$H_i = \frac{(2\pi)^4 15\pi\lambda}{U_0^7} \left(\frac{1}{x^7} + \frac{\gamma}{U_0^6} h_i \right), \quad F_1 = 1 - \frac{1}{x^7} + \frac{\gamma}{U_0^6} f_1,$$

where $i = 1, 2, 3$, and h_i and f_1 are functions of a dimensionless variable $x = \frac{U}{U_0}$. This ansatz is static and possesses SO(9) rotation symmetry, and if we take $N = \infty$, the metric just becomes the classical solution (6). By solving the equations of motion (13), we determine functions $h_i(x)$ and $f_1(x)$.

Calculations are straightforward but complicated, so we use the Mathematica code to explicitly write down the equations of motion. Some of the results are listed in the appendices A and B. From the output we find that there are five nontrivial equations which are given

by

$$E_1 = -63x^{34}f_1 - 9x^{35}f_1' - 49x^{41}h_1 + 49x^{34}(1-x^7)h_2 + 23x^{35}(1-x^7)h_2' + 2x^{36}(1-x^7)h_2'' \\ + 98x^{41}h_3 + 7x^{42}h_3' - 63402393600x^{14} + 70230343680x^7 + 1062512640 = 0, \quad (15)$$

$$E_2 = 63x^{34}f_1 + 9x^{35}f_1' + 7x^{34}(9-2x^7)h_1 + 9x^{35}(1-x^7)h_1' - 112x^{34}(1-x^7)h_2 \\ - 16x^{35}(1-x^7)h_2' - 98x^{41}h_3 - 7x^{42}h_3' - 2159861760x^7 - 5730600960 = 0, \quad (16)$$

$$E_3 = 133x^{34}f_1 + 35x^{35}f_1' + 2x^{36}f_1'' + 28x^{34}(3-10x^7)h_1 + 7x^{35}(4-7x^7)h_1' + 2x^{36}(1-x^7)h_1'' \\ - 7x^{34}(5-26x^7)h_2 - 21x^{35}(1-2x^7)h_2' - 2x^{36}(1-x^7)h_2'' + 98x^{41}h_3 + 7x^{42}h_3' \\ + 5669637120x^7 - 8626383360 = 0, \quad (17)$$

$$E_4 = 259x^{34}f_1 + 53x^{35}f_1' + 2x^{36}f_1'' + 147x^{34}(1-3x^7)h_1 + x^{35}(37-58x^7)h_1' \\ + 2x^{36}(1-x^7)h_1'' + 147x^{41}h_2 + 21x^{42}h_2' + 294x^{41}h_3 + 21x^{42}h_3' \\ - 63402393600x^{14} + 133632737280x^7 - 71292856320 = 0, \quad (18)$$

$$E_5 = 49x^{34}h_1 + 7x^{35}h_1' + 49x^{34}h_2 - x^{35}h_2' - x^{36}h_2'' - 98x^{34}h_3 - 22x^{35}h_3' - x^{36}h_3'' \\ - 63402393600x^7 + 70230343680 = 0. \quad (19)$$

Here we defined $E_1 = 4U_0^8 \ell_s^4 x^{36} \gamma^{-1} E_{00}$, $E_2 = 4U_0^8 \ell_s^4 x^{36} \gamma^{-1} E_{11}$, $E_3 = 4U_0^8 \ell_s^4 x^{36} \gamma^{-1} E_{22}$, $E_4 = 4U_0^8 \ell_s^4 x^{36} \gamma^{-1} E_{1010}$ and $E_5 = 4U_0^8 \ell_s^4 x^{\frac{65}{2}} (-1+x^7)^{-\frac{1}{2}} \gamma^{-1} E_{010}$. Note that the above equations are derived up to the order of γ , and a part of γ^0 is zero since the ansatz (14) is a fluctuation around the classical solution (6).

Now we solve these equations to obtain h_i and f_1 . We will see that h_i and f_1 are uniquely determined as functions of x by imposing reasonable boundary conditions. Because calculations below are a bit tedious, the results are summarized in the end of this section.

First let us evaluate the sum of E_1 and E_2 .

$$\frac{1}{9x^{28}(x^7-1)}(E_1 + E_2) = -7x^6h_1 - x^7h_1' + 7x^6h_2 - \frac{7}{9}x^7h_2' - \frac{2}{9}x^8h_2'' \\ + \frac{518676480}{x^{28}} - \frac{7044710400}{x^{21}} \\ = \left(-x^7h_1 + x^7h_2 - \frac{2}{9}x^8h_2' + \frac{352235520}{x^{20}} - \frac{19210240}{x^{27}} \right)' = 0. \quad (20)$$

From this equation h_1 is expressed in terms of h_2 as

$$h_1 = h_2 - \frac{2}{9}xh_2' + \frac{c_1}{x^7} + \frac{352235520}{x^{27}} - \frac{19210240}{x^{34}}, \quad (21)$$

where c_1 is an integral constant. Next let us evaluate E_5 .

$$\begin{aligned}
\frac{1}{x^{28}}E_5 &= 49x^6h_1 + 7x^7h'_1 + 49x^6h_2 - x^7h'_2 - x^8h''_2 - 98x^6h_3 - 22x^7h'_3 - x^8h''_3 \\
&\quad - \frac{63402393600}{x^{21}} + \frac{70230343680}{x^{28}} \\
&= \left(7x^7h_1 + 7x^7h_2 - x^8h'_2 - 14x^7h_3 - x^8h'_3 + \frac{3170119680}{x^{20}} - \frac{2601123840}{x^{27}}\right)' \\
&= \left(14x^7h_2 - \frac{23}{9}x^8h'_2 - 14x^7h_3 - x^8h'_3 + \frac{5635768320}{x^{20}} - \frac{2735595520}{x^{27}}\right)' = 0. \quad (22)
\end{aligned}$$

In the last line, we removed h_1 by using the eq. (21). Thus a linear combination of h_3 is expressed in terms of h_2 as

$$14x^7h_3 + x^8h'_3 = 14x^7h_2 - \frac{23}{9}x^8h'_2 + c_2 + \frac{5635768320}{x^{20}} - \frac{2735595520}{x^{27}}, \quad (23)$$

where c_2 is an integral constant. From the eqs. (21) and (23), it is possible to remove h_1 and h_3 out of E_1 , E_3 and E_4 . After some calculations, we obtain three equations remaining to be solved.

$$\begin{aligned}
E_1 &= -63x^{34}f_1 - 9x^{35}f'_1 + 49x^{34}h_2 + x^{35}(23 - 30x^7)h'_2 + 2x^{36}(1 - x^7)h''_2 \\
&\quad - 49c_1x^{34} + 7c_2x^{34} - 41211555840x^{14} + 52022476800x^7 + 1062512640 = 0, \quad (24)
\end{aligned}$$

$$\begin{aligned}
E_3 &= 133x^{34}f_1 + 35x^{35}f'_1 + 2x^{36}f''_1 \\
&\quad + 49x^{34}h_2 - \frac{7}{9}x^{35}(23 - 62x^7)h'_2 - \frac{2}{9}x^{36}(32 - 53x^7)h''_2 - \frac{4}{9}x^{37}(1 - x^7)h'''_2 \\
&\quad - 49c_1x^{34} + 7c_2x^{34} - 125748080640x^{14} + 301493283840x^7 - 37672266240 = 0, \quad (25)
\end{aligned}$$

$$\begin{aligned}
E_4 &= 259x^{34}f_1 + 53x^{35}f'_1 + 2x^{36}f''_1 \\
&\quad + 147x^{34}h_2 - \frac{7}{9}x^{35}(5 - 26x^7)h'_2 - \frac{2}{9}x^{36}(32 - 53x^7)h''_2 - \frac{4}{9}x^{37}(1 - x^7)h'''_2 \\
&\quad - 147c_1x^{34} + 21c_2x^{34} - 81366405120x^{14} + 324970168320x^7 - 95670650880. \quad (26)
\end{aligned}$$

Notice, however, that three functions E_1 , E_3 and E_4 are not independent because of the identity

$$E_4 = \frac{2}{7}xE'_1 - 9E_1 + \frac{16}{7}E_3. \quad (27)$$

This corresponds to the energy conservation, $D_a E^{ab} = 0$. Thus we only need to solve

following two equations.

$$\begin{aligned}
-\frac{1}{2}E_1 + \frac{1}{4}(E_3 - E_4) &= -\frac{1}{14}xE'_1 + \frac{7}{4}E_1 - \frac{9}{28}E_3 \\
&= -49x^{34}h_2 - x^{35}(15 - 22x^7)h'_2 - x^{36}(1 - x^7)h''_2 + 7(7c_1 - c_2)x^{34} \\
&\quad + 9510359040x^{14} - 31880459520x^7 + 13968339840 = 0, \tag{28} \\
\frac{1}{2}(E_3 - E_4) &= -\frac{1}{7}xE'_1 + \frac{9}{2}E_1 - \frac{9}{14}E_3 \\
&= -63x^{34}f_1 - 9x^{35}f'_1 - 49x^{34}h_2 - 7x^{35}(1 - 2x^7)h'_2 + 7(7c_1 - c_2)x^{34} \\
&\quad - 22190837760x^{14} - 11738442240x^7 + 28999192320 = 0. \tag{29}
\end{aligned}$$

By solving the eq. (28), finally we obtain h_2 as

$$\begin{aligned}
h_2 &= \frac{19160960}{x^{34}} - \frac{58528288}{x^{27}} + \frac{2213568}{13x^{20}} - \frac{1229760}{13x^{13}} \\
&\quad + c_1 - \frac{c_2}{7} + \frac{2459520}{x^6} + \frac{c_4}{3136x^7} + 1054080\left(2 - \frac{1}{x^7}\right)I(x), \tag{30} \\
I(x) &= \frac{c_3}{944455680} + \log(x - 1) + \frac{c_4}{6611189760} \log(1 - x^{-7}) \\
&\quad - \sum_{n=1,3,5} \cos \frac{n\pi}{7} \log(x^2 + 2x \cos \frac{n\pi}{7} + 1) \\
&\quad - 2 \sum_{n=1,3,5} \sin \frac{n\pi}{7} \tan^{-1} \left(\frac{x + \cos \frac{n\pi}{7}}{\sin \frac{n\pi}{7}} \right), \tag{31}
\end{aligned}$$

where c_3 and c_4 are integral constants. Although the form of $I(x)$ seems to be complicated, its derivative becomes

$$I'(x) = \frac{7}{x^7 - 1} \left(1 + \frac{c_4 x^{-1}}{6611189760} \right). \tag{32}$$

So far there are four integral constants, but these will be fixed by appropriate conditions. In fact it is natural to require that $h_i(1)$ are finite and $h_i(x) \sim \mathcal{O}(x^{-8})$ when x goes to the infinity. In order to satisfy these conditions, it is necessary to choose $c_2 = 7c_1$, $c_3 = 944455680\pi(\sin \frac{\pi}{7} + \sin \frac{3\pi}{7} + \sin \frac{5\pi}{7})$ and $c_4 = -6611189760$. Inserting these values into the eqs. (30), (31) and (32), we obtain

$$\begin{aligned}
h_2 &= \frac{19160960}{x^{34}} - \frac{58528288}{x^{27}} + \frac{2213568}{13x^{20}} - \frac{1229760}{13x^{13}} \\
&\quad - \frac{2108160}{x^7} + \frac{2459520}{x^6} + 1054080\left(2 - \frac{1}{x^7}\right)I(x), \tag{33} \\
I(x) &= \log \frac{x^7(x-1)}{x^7-1} - \sum_{n=1,3,5} \cos \frac{n\pi}{7} \log(x^2 + 2x \cos \frac{n\pi}{7} + 1) \\
&\quad - 2 \sum_{n=1,3,5} \sin \frac{n\pi}{7} \left\{ \tan^{-1} \left(\frac{x + \cos \frac{n\pi}{7}}{\sin \frac{n\pi}{7}} \right) - \frac{\pi}{2} \right\}, \tag{34}
\end{aligned}$$

and

$$I'(x) = \frac{7(1-x^{-1})}{x^7-1}. \quad (35)$$

Note that the function $I(x)$ behaves as

$$I(x) \sim -\frac{7}{6x^6} + \frac{1}{x^7} - \frac{7}{13x^{13}} + \frac{1}{2x^{14}} + \mathcal{O}(x^{-15}), \quad (36)$$

when x goes to the infinity.

Now we remove h_2 out of the eq. (29), and obtain the differential equation only for f_1 .

$$\begin{aligned} \frac{1}{18x^{28}}(E_3 - E_4) &= -x^7 f_1' - 7x^6 f_1 + 819840I' + 3279360x^7(x^7-1)I' \\ &+ \frac{3624512640}{x^{28}} - \frac{3228113280}{x^{21}} - \frac{5738880}{x^{14}} - \frac{5738880}{x^7} \\ &+ 22955520x^6 - 22955520x^7 \\ &= \left(-x^7 f_1 + 819840I - \frac{1208170880}{9x^{27}} \right. \\ &\quad \left. + \frac{161405664}{x^{20}} + \frac{5738880}{13x^{13}} + \frac{956480}{x^6} \right)' = 0. \end{aligned} \quad (37)$$

Then f_1 is solved as

$$f_1 = -\frac{1208170880}{9x^{34}} + \frac{161405664}{x^{27}} + \frac{5738880}{13x^{20}} + \frac{956480}{x^{13}} + \frac{819840}{x^7} I(x). \quad (38)$$

Here the integral constant is set to be zero, because we imposed the boundary condition that $f_1(x) \sim \mathcal{O}(x^{-8})$ when x goes to the infinity. From the eq. (21), h_1 is determined as

$$\begin{aligned} h_1 &= \frac{1302501760}{9x^{34}} - \frac{57462496}{x^{27}} + \frac{12051648}{13x^{20}} - \frac{4782400}{13x^{13}} \\ &- \frac{3747840}{x^7} + \frac{4099200}{x^6} - \frac{1639680(x-1)}{(x^7-1)} + 117120 \left(18 - \frac{23}{x^7} \right) I(x). \end{aligned} \quad (39)$$

The integral constant c_1 is chosen to be zero so as to satisfy $h_1(x) \sim \mathcal{O}(x^{-8})$ when x goes to the infinity. Finally from the eq. (23), we derive

$$\begin{aligned} 0 &= -x^{14}h_3' - 14x^{13}h_3 + (29514240x^{13} - 33613440x^6)I(x) \\ &+ (2693760x^7 - 5387520x^{14})I'(x) + 72145920x^7 - 67226880x^6 \\ &- \frac{7222208000}{9x^{21}} + \frac{777920416}{x^{14}} + \frac{144127872}{13x^7} - \frac{58072000}{13} \\ &= \left(-x^{14}h_3 + (2108160x^{14} - 4801920x^7)I(x) + 2459520x^8 - 2108160x^7 \right. \\ &\quad \left. + \frac{361110400}{9x^{20}} - \frac{59840032}{x^{13}} - \frac{24021312}{13x^6} - \frac{58072000}{13}x \right)'. \end{aligned} \quad (40)$$

Thus h_3 is expressed as

$$\begin{aligned} h_3 &= \frac{361110400}{9x^{34}} - \frac{59840032}{x^{27}} - \frac{24021312}{13x^{20}} - \frac{58072000}{13x^{13}} \\ &- \frac{2108160}{x^7} + \frac{2459520}{x^6} + 117120 \left(18 - \frac{41}{x^7} \right) I(x). \end{aligned} \quad (41)$$

The integral constant is set to be zero, since this term can be removed by the general coordinate transformation on z direction. It corresponds to the gauge transformation on C_μ in ten dimensions.

Let us summarize the quantum correction to the near horizon geometry of the non-extremal M-wave and the black 0-brane. By solving the eqs. (15)–(19), we obtained the quantum near horizon geometry of the non-extremal M-wave,

$$ds_{11}^2 = \ell_s^4 \left(-H_1^{-1} F_1 dt^2 + F_1^{-1} U_0^2 dx^2 + U_0^2 x^2 d\Omega_8^2 + (\ell_s^{-4} H_2^{\frac{1}{2}} dz - H_3^{-\frac{1}{2}} dt)^2 \right), \quad (42)$$

$$H_i = \frac{(2\pi)^4 15\pi\lambda}{U_0^7} \left(\frac{1}{x^7} + \epsilon \frac{\lambda^2}{U_0^6} h_i \right), \quad F_1 = 1 - \frac{1}{x^7} + \epsilon \frac{\lambda^2}{U_0^6} f_1.$$

In stead of γ , we introduced dimensionless parameter

$$\epsilon = \frac{\gamma}{\lambda^2} = \frac{\pi^6}{2^7 3^2 N^2} \sim \frac{0.835}{N^2}, \quad (43)$$

and the functions h_i and f_1 are uniquely determined as

$$\begin{aligned} h_1 &= \frac{1302501760}{9x^{34}} - \frac{57462496}{x^{27}} + \frac{12051648}{13x^{20}} - \frac{4782400}{13x^{13}} \\ &\quad - \frac{3747840}{x^7} + \frac{4099200}{x^6} - \frac{1639680(x-1)}{(x^7-1)} + 117120 \left(18 - \frac{23}{x^7} \right) I(x), \\ h_2 &= \frac{19160960}{x^{34}} - \frac{58528288}{x^{27}} + \frac{2213568}{13x^{20}} - \frac{1229760}{13x^{13}} \\ &\quad - \frac{2108160}{x^7} + \frac{2459520}{x^6} + 1054080 \left(2 - \frac{1}{x^7} \right) I(x), \\ h_3 &= \frac{361110400}{9x^{34}} - \frac{59840032}{x^{27}} - \frac{24021312}{13x^{20}} - \frac{58072000}{13x^{13}} \\ &\quad - \frac{2108160}{x^7} + \frac{2459520}{x^6} + 117120 \left(18 - \frac{41}{x^7} \right) I(x), \\ f_1 &= -\frac{1208170880}{9x^{34}} + \frac{161405664}{x^{27}} + \frac{5738880}{13x^{20}} + \frac{956480}{x^{13}} + \frac{819840}{x^7} I(x). \end{aligned} \quad (44)$$

The function $I(x)$ is defined by the eq. (34). In order to fix the integral constants, we required that $h_i(1)$ are finite and $h_i(x), f_1(x) \sim \mathcal{O}(x^{-8})$ when x goes to the infinity. After the dimensional reduction to ten dimensions, we obtain

$$ds_{10}^2 = \ell_s^2 \left(-H_1^{-1} H_2^{\frac{1}{2}} F_1 dt^2 + H_2^{\frac{1}{2}} F_1^{-1} U_0^2 dx^2 + H_2^{\frac{1}{2}} U_0^2 x^2 d\Omega_8^2 \right), \quad (45)$$

$$e^\phi = \ell_s^{-3} H_2^{\frac{3}{4}}, \quad C = \ell_s^4 H_2^{-\frac{1}{2}} H_3^{-\frac{1}{2}} dt.$$

This represents the quantum near horizon geometry of the non-extremal black 0-brane.

5 Thermodynamics of Quantum Near Horizon Geometry of Black 0-Brane

Since the quantum near horizon geometry of the non-extremal black 0-brane is derived in the previous section, it is interesting to evaluate its thermodynamics. In this section, we estimate the entropy and the internal energy of the quantum near horizon geometry of the non-extremal black 0-brane by using Wald's formula [41, 42]. These quantities are quite important when we test the gauge/gravity duality.

In the following, quantities are calculated up to $\mathcal{O}(\epsilon^2)$. First of all, let us examine the location of the horizon x_H . This is defined by $F_1(x_H) = 0$ and becomes

$$x_H = 1 - \epsilon \frac{f_1(1)}{7} \tilde{U}_0^{-6}, \quad (46)$$

where $\tilde{U}_0 \equiv U_0/\lambda^{\frac{1}{3}}$ is a dimensionless parameter. Temperature of the black 0-brane is derived by the usual prescription. We consider the Euclidean geometry by changing time coordinate as $t = -i\tau$ and require the smoothness of the geometry at the horizon. This fixes the periodicity of τ direction and its inverse gives the temperature of the non-extremal black 0-brane. Then the dimensionless temperature $\tilde{T} = T/\lambda^{\frac{1}{3}}$ of the black 0-brane is evaluated as

$$\tilde{T} = \frac{1}{4\pi} U_0^{-1} H_1^{-\frac{1}{2}} F_1' \Big|_{x_H} / \lambda^{\frac{1}{3}} = a_1 \tilde{U}_0^{\frac{5}{2}} (1 + \epsilon a_2 \tilde{U}_0^{-6}), \quad (47)$$

where a_1 and a_2 are numerical constants given by

$$\begin{aligned} a_1 &= \frac{7}{16\pi^3 \sqrt{15}\pi} \sim 0.00206, \\ a_2 &= \frac{9}{14} f_1(1) + \frac{1}{7} f_1'(1) - \frac{1}{2} h_1(1) \sim 937000. \end{aligned} \quad (48)$$

Inversely solving the eq. (47), the dimensionless parameter \tilde{U}_0 is written in terms of the temperature \tilde{T} as

$$\tilde{U}_0 = a_1^{-\frac{2}{5}} \tilde{T}^{\frac{2}{5}} \left(1 - \epsilon \frac{2}{5} a_1^{\frac{12}{5}} a_2 \tilde{T}^{-\frac{12}{5}} \right), \quad (49)$$

By using this replacement, it is always possible to express physical quantities as functions of \tilde{T} .

Next we derive the entropy of the quantum near horizon geometry of the non-extremal black 0-brane. In practice, we consider the quantum near horizon geometry of the non-extremal M-wave because of its simple expression. Since the effective action (9) includes higher derivative terms, we should employ Wald's entropy formula which ensures the first law of the black hole thermodynamics. The Wald's entropy formula is given by

$$S = -2\pi \int_H d\Omega_8 dz \sqrt{h} \frac{\partial S_{11}}{\partial R_{\mu\nu\rho\sigma}} N_{\mu\nu} N_{\rho\sigma}, \quad (50)$$

where $\sqrt{h} = (\ell_s^2 U_0 x)^8 \ell_s^{-2} H_2^{1/2}$ is the volume factor at the horizon and $N_{\mu\nu}$ is an antisymmetric tensor binormal to the horizon. The binormal tensor satisfies $N_{\mu\nu} N^{\mu\nu} = -2$ and nonzero component is only $N_{tx} = -\ell_s^4 U_0 H_1^{-1/2}$. The effective action is given by the eq. (9), and in the formula the variation of the action is evaluated as if the Riemann tensor is an independent variable, that is,

$$\frac{\partial S_{11}}{\partial R_{\mu\nu\rho\sigma}} = \frac{1}{2\kappa_{11}^2} (g^{\mu[\rho} g^{\sigma]\nu} + \gamma \ell_s^{12} X^{\mu\nu\rho\sigma}). \quad (51)$$

Now we are ready to evaluate the entropy of the quantum near horizon geometry of the non-extremal M-wave. Some useful results are collected in the appendix B. By using these, the entropy is evaluated as

$$\begin{aligned} S &= \frac{4\pi}{2\kappa_{11}^2} \int_{\text{H}} d\Omega_8 dz \sqrt{h} \left(1 - \frac{1}{2} \gamma \ell_s^{12} X^{\mu\nu\rho\sigma} N_{\mu\nu} N_{\rho\sigma} \right) \\ &= \frac{4\pi}{2\kappa_{11}^2} \int_{\text{H}} d\Omega_8 dz \sqrt{h} (1 - 2\gamma \ell_s^{20} U_0^2 H_1^{-1} X^{txtx}) \\ &= \frac{4\pi}{2\kappa_{11}^2} \int_{\text{H}} d\Omega_8 dz \sqrt{h} \left(1 + 40642560 \epsilon \frac{1}{\tilde{U}_0^6 x_{\text{H}}^{20}} \right) \\ &= \frac{4}{49} a_1 N^2 \tilde{U}_0^{\frac{9}{2}} \left\{ 1 + \epsilon \left(-\frac{9}{14} f_1(1) + \frac{1}{2} h_2(1) + 40642560 \right) \tilde{U}_0^{-6} \right\} \\ &= \frac{4}{49} a_1^{-\frac{4}{5}} N^2 \tilde{T}^{\frac{9}{5}} \left\{ 1 + \epsilon a_1^{\frac{12}{5}} \left(-\frac{9}{5} f_1(1) - \frac{9}{35} f_1'(1) + \frac{9}{10} h_1(1) + \frac{1}{2} h_2(1) + 40642560 \right) \tilde{T}^{-\frac{12}{5}} \right\} \\ &= a_3 N^2 \tilde{T}^{\frac{9}{5}} (1 + \epsilon a_4 \tilde{T}^{-\frac{12}{5}}), \end{aligned} \quad (52)$$

where numerical constants a_3 and a_4 are defined as

$$\begin{aligned} a_3 &= \frac{4}{49} a_1^{-\frac{4}{5}} = 2^{\frac{26}{5}} 15^{\frac{2}{5}} 7^{-\frac{14}{5}} \pi^{\frac{14}{5}} \sim 11.5, \\ a_4 &= a_1^{\frac{12}{5}} \left(-\frac{9}{5} f_1(1) - \frac{9}{35} f_1'(1) + \frac{9}{10} h_1(1) + \frac{1}{2} h_2(1) + 40642560 \right) \sim 0.400. \end{aligned} \quad (53)$$

So far we have obtained the entropy for the M-wave. Because of the duality between type IIA string theory and M-theory, this is equivalent to that of the black 0-brane.

Finally let us derive the internal energy of the quantum near horizon geometry of the non-extremal black 0-brane. Wald's entropy formula is constructed so as to satisfy the thermodynamic laws of black holes. Then by integrating $d\tilde{E} = \tilde{T} dS$, it is possible to obtain the dimensionless energy $\tilde{E} = E/\lambda^{\frac{1}{3}}$ as

$$\frac{\tilde{E}}{N^2} = \frac{9}{14} a_3 \tilde{T}^{\frac{14}{5}} - \epsilon \frac{3}{2} a_3 a_4 \tilde{T}^{\frac{2}{5}} \sim 7.41 \tilde{T}^{\frac{14}{5}} - \frac{5.77}{N^2} \tilde{T}^{\frac{2}{5}}. \quad (54)$$

This result includes the quantum gravity effect, and it gives quite nontrivial test of the gauge/gravity duality if we can evaluate the internal energy from the dual gauge theory. In fact it is possible by employing the Monte Carlo simulation and the result strongly concludes that the duality holds at this order [24].

The specific heat is evaluated as

$$\frac{1}{N^2} \frac{d\tilde{E}}{d\tilde{T}} = \frac{9}{5} a_3 \tilde{T}^{\frac{9}{5}} - \epsilon \frac{3}{5} a_3 a_4 \tilde{T}^{-\frac{3}{5}}. \quad (55)$$

Notice that the specific heat becomes negative in the region where $\tilde{T} < (\epsilon a_4/3)^{5/12} \sim 0.4N^{-5/6}$. In this region the non-extremal black 0-brane behaves like Schwarzschild black hole and will be unstable. When $N = \infty$ the instability will be suppressed. This result is also verified from the Monte Carlo simulation of the dual gauge theory [24].

6 D0-brane Probe

In this section, we probe the quantum near horizon geometry of the non-extremal black 0-brane (45) via a D0-brane. From the analysis it is possible to study how the test D0-brane is affected by the background field.

The bosonic part of the D0-brane action consists of the Born-Infeld action and the Chern-Simons one. Here we neglect an excitation of the gauge field on the D0-brane, so the Born-Infeld action is simply given by the pull-back of the metric. We also assume that the D0-brane moves only along the radial direction. Then the probe D0-brane action in the background of (45) is written as

$$\begin{aligned} S_{\text{D0}} &= -T_0 \int dt e^{-\phi} \sqrt{-g_{\mu\nu} \frac{dx^\mu}{dt} \frac{dx^\nu}{dt}} + T_0 \int C \\ &= -T_0 \ell_s^4 \int dt H_2^{-\frac{1}{2}} \sqrt{H_1^{-1} F_1 - F_1^{-1} U_0^2 \dot{x}^2} + T_0 \ell_s^4 \int dt H_2^{-\frac{1}{2}} H_3^{-\frac{1}{2}}. \end{aligned} \quad (56)$$

The momentum conjugate to x is evaluated as

$$p = T_0 \ell_s^4 H_2^{-\frac{1}{2}} \frac{F_1^{-1} U_0^2 \dot{x}}{\sqrt{H_1^{-1} F_1 - F_1^{-1} U_0^2 \dot{x}^2}}, \quad (57)$$

and the energy of the probe D0-brane is given by

$$\begin{aligned} E_{\text{D0}} &= p\dot{x} + T_0 \ell_s^4 H_2^{-\frac{1}{2}} \sqrt{H_1^{-1} F_1 - F_1^{-1} U_0^2 \dot{x}^2} - T_0 \ell_s^4 H_2^{-\frac{1}{2}} H_3^{-\frac{1}{2}} \\ &= T_0 \ell_s^4 H_2^{-\frac{1}{2}} \frac{H_1^{-1} F_1}{\sqrt{H_1^{-1} F_1 - F_1^{-1} U_0^2 \dot{x}^2}} - T_0 \ell_s^4 H_2^{-\frac{1}{2}} H_3^{-\frac{1}{2}} \\ &= T_0 \ell_s^4 H_1^{-\frac{1}{2}} H_2^{-\frac{1}{2}} F_1^{\frac{1}{2}} \sqrt{1 + \left(\frac{p F_1^{\frac{1}{2}} H_2^{\frac{1}{2}}}{T_0 \ell_s^4 U_0} \right)^2} - T_0 \ell_s^4 H_2^{-\frac{1}{2}} H_3^{-\frac{1}{2}} \\ &\sim \frac{1}{2} H_1^{-\frac{1}{2}} H_2^{\frac{1}{2}} F_1^{\frac{3}{2}} \frac{p^2}{T_0 \ell_s^4 U_0^2} + T_0 \ell_s^4 \left(H_1^{-\frac{1}{2}} H_2^{-\frac{1}{2}} F_1^{\frac{1}{2}} - H_2^{-\frac{1}{2}} H_3^{-\frac{1}{2}} \right). \end{aligned} \quad (58)$$

In the final line we took the non-relativistic limit. From this we see that the potential energy for the probe D0-brane is expressed as

$$V_{D0} = T_0 \ell_s^4 \left(H_1^{-\frac{1}{2}} H_2^{-\frac{1}{2}} F_1^{\frac{1}{2}} - H_2^{-\frac{1}{2}} H_3^{-\frac{1}{2}} \right). \quad (59)$$

The first term corresponds to the gravitational attractive force and the second one does to the R-R repulsive force.

When we take $N = \infty$, the potential energy becomes $V_{D0} = T_0 \ell_s^4 H^{-1} (\sqrt{F} - 1)$. The part $(\sqrt{F} - 1)$ shows that the gravitational attractive force overcomes the R-R repulsive force. Similarly, when N is finite, we regard $\sqrt{F_1}$ as the gravitational attractive force to the probe D0-brane. The function of $\sqrt{F_1}$ is plotted in fig. 1. From this we see that the gravitational force becomes repulsive near the horizon x_H .




Figure 1: The function $\sqrt{F_1(x)}$ with $F_1(x) = 1 - 1/x^7 + 0.000001f_1(x)$.

7 Conclusion and Discussion

In this paper we studied quantum nature of the bunch of D0-branes in the type IIA superstring theory. In the classical limit, it is well described by the non-extremal black 0-brane in the type IIA supergravity. The quantum correction to the non-extremal black 0-brane is investigated after taking the near horizon limit.

In order to manage the quantum effect of the gravity, we uplifted the near horizon geometry of the non-extremal black 0-brane into that of the M-wave solution in the eleven dimensional supergravity. These two are equivalent via the duality between the type IIA

superstring theory and the M-theory, but the latter is purely geometrical and calculations become rather simple. The geometrical part of the effective action for the M-theory (9) is derived so as to be consistent with the 1-loop amplitudes in the type IIA superstring theory. And the quantum correction to the M-wave solution is taken into account by explicitly solving the equations of motion (13). The solution is uniquely determined and its explicit form is given by the eq. (45). It is interesting to note that a probe D0-brane moving in this background would feel repulsive force near the horizon. It means that the solution includes the back-reaction of the Hawking radiation.

We also investigated the thermodynamic property of the quantum near horizon geometry of the non-extremal black 0-brane. Since the effective action contains higher derivative terms, we examined the thermodynamic property of the black 0-brane by employing Wald's formula. The entropy and the internal energy of the black 0-brane are evaluated up to $1/N^2$. The quantum correction to the internal energy becomes important when N is small. In ref. [24], the internal energy is also calculated from the dual thermal gauge theory by using the Monte Carlo simulation, and it agrees with the eq. (54) very well. This gives a strong evidence for the gauge/gravity duality at the level of quantum gravity.

Finally we give an important remark on the effective action for the M-theory. It contains higher derivative terms, but these cannot be determined uniquely because of the field redefinitions. In the appendices we have considered all possible higher derivative terms and shown that the ambiguities of the effective action have nothing to do with the thermodynamic properties of near horizon geometry of the non-extremal black 0-brane.

As a future work, it is important to derive quantum geometry of the non-extremal black 0-brane and obtain the solution (45) by taking the near horizon limit. The result will be reported elsewhere, but it is really possible. It is also interesting to examine quantum correction to the black 6-brane, which is also described by purely geometrical object, called Kaluza-Klein monopole, in the eleven dimensional supergravity. To find connections of our results to the other approaches to the field theory on the D0-branes is important as well [43, 44]. Since now we capture the quantum nature of the near horizon geometry of the black 0-brane, it is interesting to consider a recent proposal to resolve the information paradox on the black hole [45, 46].

Acknowledgement

The author would like to thank Masanori Hanada, Goro Ishiki and Jun Nishimura for discussions and collaborations. He would also like to thank Masafumi Fukuma, Hideki Ishihara, Hikaru Kawai and Yukinori Yasui for discussions. This work was partially supported by the

Ministry of Education, Science, Sports and Culture, Grant-in-Aid for Young Scientists (B)
19740141, 2007 and 24740140, 2012.

A Calculations of Ricci Tensor and Scalar Curvature

By using the ansatz (14) for the metric, each component of the Ricci tensor up to the linear order of γ is calculated as

$$\begin{aligned}
R_{00} &= \frac{\gamma}{4U_0^8 x^2 \ell_s^4} \{ 98f_1 + 30xf_1' + 2x^2 f_1'' + 49(2 - 7x^7)h_1 + 3x(10 - 17x^7)h_1' + 2x^2(1 - x^7)h_1'' \\
&\quad + 147x^7 h_2 + 21x^8 h_2' + 196x^7 h_3 + 14x^8 h_3' \}, \\
R_{11} &= \frac{\gamma}{4U_0^8 x^2 \ell_s^4} \{ -98f_1 - 30xf_1' - 2x^2 f_1'' - 35(1 - 8x^7)h_1 - 21x(1 - 2x^7)h_1' - 2x^2(1 - x^7)h_1'' \\
&\quad - 7(9 + 12x^7)h_2 + 7x(1 - 4x^7)h_2' + 2x^2(1 - x^7)h_2'' - 196x^7 h_3 - 14x^8 h_3' \}, \quad (60) \\
R_{\bar{a}\bar{a}} &= \frac{\gamma}{2U_0^8 x^2 \ell_s^4} \{ -14f_1 - 2xf_1' - 7(1 - x^7)h_1 - x(1 - x^7)h_1' + 7(1 - x^7)h_2 + x(1 - x^7)h_2' \}, \\
R_{\natural\natural} &= \frac{\gamma}{4U_0^8 x^2 \ell_s^4} \{ 98f_1 + 14xf_1' + 49(1 - 3x^7)h_1 + 7x(1 - x^7)h_1' \\
&\quad + 49(1 - x^7)h_2 + 23x(1 - x^7)h_2' + 2x^2(1 - x^7)h_2'' + 196x^7 h_3 + 14x^8 h_3' \}, \\
R_{0\natural} &= \frac{\gamma x^{3/2} \sqrt{x^7 - 1}}{4U_0^8 \ell_s^4} \{ 49h_1 + 7xh_1' + 49h_2 - xh_2' - x^2 h_2'' - 98h_3 - 22xh_3' - x^2 h_3'' \}.
\end{aligned}$$

Here we used \natural instead of 10 and $\bar{a} = 2, \dots, 9$. Ricci scalar up to the linear order of γ becomes like

$$\begin{aligned}
R &= \frac{\gamma}{2U_0^8 x^2 \ell_s^4} \{ -161f_1 - 39xf_1' - 2x^2 f_1'' - 98(1 - 3x^7)h_1 - 3x(10 - 17x^7)h_1' - 2x^2(1 - x^7)h_1'' \\
&\quad + 49(1 - 4x^7)h_2 + x(23 - 44x^7)h_2' + 2x^2(1 - x^7)h_2'' - 98x^7 h_3 - 7x^8 h_3' \}. \quad (61)
\end{aligned}$$

B Calculations of Higher Derivative Terms

In this appendix we summarize the values of higher derivative terms appeared in the eq. (13). Note that we only need to evaluate these terms by using the ansatz (14) with $\gamma = 0$, because the equations of motion are solved up to the linear order of γ . First of all, each component of R_{abcd} is calculated as

$$\begin{aligned}
R_{0101} &= -\frac{28}{U_0^2 x^2 \ell_s^4}, & R_{0\bar{a}0\bar{a}} &= \frac{7}{2U_0^2 x^2 \ell_s^4}, \\
R_{011\natural} &= \frac{28\sqrt{x^7 - 1}}{U_0^2 x^{\frac{1}{2}} \ell_s^4}, & R_{0\bar{a}\bar{a}\natural} &= -\frac{7\sqrt{x^7 - 1}}{2U_0^2 x^{\frac{1}{2}} \ell_s^4}, \\
R_{1\natural1\natural} &= -\frac{28(x^7 - 1)}{U_0^2 x^9 \ell_s^4}, & R_{1\bar{a}1\bar{a}} &= -\frac{7}{2U_0^2 x^9 \ell_s^4}, \\
R_{\bar{a}\natural\bar{a}\natural} &= \frac{7(x^7 - 1)}{2U_0^2 x^9 \ell_s^4}, & R_{\bar{a}\bar{a}\bar{a}\bar{a}} &= \frac{1}{U_0^2 x^9 \ell_s^4}.
\end{aligned} \quad (62)$$

We used \natural instead of 10 and $\bar{a}, \bar{b} = 2, \dots, 9$. The scalar curvature and each component of the Ricci tensor become zero, and each component of X_{abcd} in the eq. (12) is evaluated as

$$\begin{aligned}
X_{0101} &= -\frac{20321280}{U_0^6 x^{20} \ell_s^{12}}, & X_{0\bar{a}0\bar{a}} &= -\frac{1270080}{U_0^6 x^{20} \ell_s^{12}}, \\
X_{011\natural} &= \frac{20321280\sqrt{x^7-1}}{U_0^6 x^{\frac{47}{2}} \ell_s^{12}}, & X_{0\bar{a}\bar{a}\natural} &= \frac{1270080\sqrt{x^7-1}}{U_0^6 x^{\frac{47}{2}} \ell_s^{12}}, \\
X_{1\natural1\natural} &= -\frac{20321280(x^7-1)}{U_0^6 x^{27} \ell_s^{12}}, & X_{1\bar{a}1\bar{a}} &= \frac{1270080}{U_0^6 x^{27} \ell_s^{12}}, \\
X_{\bar{a}\natural\bar{a}\natural} &= -\frac{1270080(x^7-1)}{U_0^6 x^{27} \ell_s^{12}}, & X_{\bar{a}\bar{b}\bar{a}\bar{b}} &= \frac{1192320}{U_0^6 x^{27} \ell_s^{12}}.
\end{aligned} \tag{63}$$

By using these results we are ready to calculate higher derivative terms in the eq. (13). The R^4 terms are calculated as

$$t_8 t_8 R^4 - \frac{1}{4!} \epsilon_{11} \epsilon_{11} R^4 = \frac{531256320}{U_0^8 x^{36} \ell_s^{16}}. \tag{64}$$

The RX terms become

$$\begin{aligned}
R_{abc0} X^{abc}{}_0 &= -\frac{1066867200}{U_0^8 x^{29} \ell_s^{16}}, & R_{abc1} X^{abc}{}_1 &= \frac{1066867200}{U_0^8 x^{36} \ell_s^{16}}, \\
R_{abc\natural} X^{abc}{}_{\natural} &= -\frac{1066867200(x^7-1)}{U_0^8 x^{36} \ell_s^{16}}, & R_{abc\bar{a}} X^{abc}{}_{\bar{b}} &= -\frac{1088640}{U_0^8 x^{36} \ell_s^{16}} \delta_{\bar{a}\bar{b}}, \\
R_{abc0} X^{abc}{}_{\natural} &= R_{abc\natural} X^{abc}{}_0 = -\frac{1066867200\sqrt{x^7-1}}{U_0^8 x^{\frac{65}{2}} \ell_s^{16}},
\end{aligned} \tag{65}$$

and the DDX terms are evaluated as

$$\begin{aligned}
D_{(a} D_b) X^a{}_{00}{}^b &= \frac{198132480(-47+40x^7)}{U_0^8 x^{29} \ell_s^{16}}, & D_{(a} D_b) X^a{}_{11}{}^b &= \frac{2177280(513+124x^7)}{U_0^8 x^{36} \ell_s^{16}}, \\
D_{(a} D_b) X^a{}_{\natural\natural}{}^b &= \frac{198132480(47-87x^7+40x^{14})}{U_0^8 x^{36} \ell_s^{16}}, & D_{(a} D_b) X^a{}_{\bar{a}\bar{b}}{}^b &= \frac{236234880(4-3x^7)}{U_0^8 x^{36} \ell_s^{16}} \delta_{\bar{a}\bar{b}}, \\
D_{(a} D_b) X^a{}_{0\natural}{}^b &= D_{(a} D_b) X^a{}_{\natural 0}{}^b = \frac{198132480(-47+40x^7)\sqrt{x^7-1}}{U_0^8 x^{\frac{65}{2}} \ell_s^{16}}.
\end{aligned} \tag{66}$$

By inserting these results into the eq. (13), we obtain the eqs. (15)–(19).

C Generic R^4 Terms, Equations of Motion and Solution

In this appendix, we classify independent R4 terms which consist of four products of the Riemann tensor, the Ricci tensor or the scalar curvature. The R4 terms which include the Ricci tensor or the scalar curvature cannot be determined from the scattering amplitudes in the type IIA superstring theory. So in general the effective action and equations of motion are affected by these ambiguities.

First let us review the R4 terms which only consist of the Riemann tensor. Since there are 16 indices, we have 8 pairs to be contracted. Naively it seems that there are so many possible patterns. However, carefully using properties of the Riemann tensor, such as $R_{abcd} = -R_{bcad} - R_{cabd}$, it is possible to show that there are only 7 independent terms.

$$\begin{aligned}
B_1 &= R_{abcd}R_{abcd}R_{efgh}R_{efgh}, & B_2 &= R_{abcd}R_{aefg}R_{bcdh}R_{efgh}, \\
B_3 &= R_{abcd}R_{abef}R_{cdgh}R_{efgh}, & B_4 &= R_{acbd}R_{aebf}R_{cgdh}R_{efgh}, \\
B_5 &= R_{abcd}R_{aefg}R_{befh}R_{cdgh}, & B_6 &= R_{abcd}R_{aefg}R_{bfch}R_{cdgh}, \\
B_7 &= R_{acbd}R_{aefg}R_{befh}R_{cgdh}.
\end{aligned} \tag{67}$$

In the main part of this paper we considered the R4 terms $t_8 t_8 R^4 - \frac{1}{4!} \epsilon_{11} \epsilon_{11} R^4 = 24(B_1 - 64B_2 + 2B_3 + 16B_4 - 16B_5 - 16B_6)$ which is explicitly written in the eq. (8). In order to derive equations of motion, we need to calculate variations of (67). These are evaluated as

$$\begin{aligned}
\delta B_1 &= 4(\delta R_{abcd})R_{abcd}R_{efgh}R_{efgh}, & \delta B_2 &= (\delta R_{abcd})R_{abce}R_{dfgh}R_{efgh}, \\
\delta B_3 &= 4(\delta R_{abcd})R_{abef}R_{cdgh}R_{efgh}, & \delta B_4 &= 4(\delta R_{abcd})R_{aecg}R_{bfch}R_{efgh}, \\
\delta B_5 &= 2(\delta R_{abcd})R_{abeg}R_{cfch}R_{dfgh} + 2(\delta R_{abcd})R_{efag}R_{efch}R_{gbhd}, \\
\delta B_6 &= 2(\delta R_{abcd})R_{abeg}R_{cfch}R_{dfgh} + 2(\delta R_{abcd})R_{efag}R_{efch}R_{gbhd} - 2(\delta R_{abcd})R_{abef}R_{cegh}R_{dfgh}, \\
\delta B_7 &= 4(\delta R_{abcd})R_{aefg}R_{cefh}R_{gbhd}.
\end{aligned} \tag{68}$$

By using these results, we evaluated the eq. (11) and derived the equations of motion (13).

Next let us consider the R4 terms which necessarily depend on the Ricci tensor or the scalar curvature. Since the procedure for the classification is straightforward, we employ a Mathematica code. As a result those are classified into 19 terms.

$$\begin{aligned}
B_8 &= R_{abcd}R_{abcd}R_{ef}R_{ef}, & B_9 &= R_{abcd}R_{abcd}R^2, & B_{10} &= R_{abcd}R_{bcd}R_{ef}R_{ae}, \\
B_{11} &= R_{abcd}R_{aefg}R_{bcdg}R_{ef}, & B_{12} &= R_{abcd}R_{bcde}R_{ae}R, & B_{13} &= R_{acbd}R_{cedf}R_{ef}R_{ab}, \\
B_{14} &= R_{abcd}R_{abeg}R_{cdfg}R_{ef}, & B_{15} &= R_{acbd}R_{aebg}R_{cfdg}R_{ef}, & B_{16} &= R_{abcd}R_{abef}R_{cdef}R, \\
B_{17} &= R_{acbd}R_{aebf}R_{cedf}R, & B_{18} &= R_{acbd}R_{ab}R_{cd}R, & B_{19} &= R_{abcd}R_{cdef}R_{ae}R_{bf}, \\
B_{20} &= R_{acbd}R_{cedf}R_{ae}R_{bf}, & B_{21} &= R_{acbd}R_{ae}R_{be}R_{cd}, & B_{22} &= R_{ab}R_{ab}R_{cd}R_{cd}, \\
B_{23} &= R_{ab}R_{ab}R^2, & B_{24} &= R_{ab}R_{cd}R_{ac}R_{bd}, & B_{25} &= R_{ab}R_{ac}R_{bc}R, \\
B_{26} &= R^4.
\end{aligned} \tag{69}$$

Then the effective action (9) is generalized into the form of

$$S_{11} = \frac{1}{2\kappa_{11}^2} \int d^{11}x e \left\{ R + \gamma \ell_s^{12} \left(t_8 t_8 R^4 - \frac{1}{4!} \epsilon_{11} \epsilon_{11} R^4 + \sum_{n=8}^{26} b_n B_n \right) \right\}. \tag{70}$$

The coefficients $b_n (n = 8, \dots, 26)$ cannot be determined from the results of scattering amplitudes in the type IIA superstring theory, since we can remove or add these terms by appropriate field redefinitions of the metric. Therefore it is expected that these terms do not affect physical quantities such as the internal energy of the black 0-brane. We will confirm this in the appendix D.

Let us derive equations of motion for the effective action (70). The variations of 19 terms in (69) are evaluated as

$$\begin{aligned}
\delta B_8 &= (\delta R_{abcd})(2R_{abcd}R_{ef}R_{ef} + 2R_{efgh}R_{efgh}R_{ac}\eta_{bd}), \\
\delta B_9 &= (\delta R_{abcd})(2R_{abcd}R^2 + 2R_{efgh}R_{efgh}\eta_{ac}\eta_{bd}R), \\
\delta B_{10} &= (\delta R_{abcd})(R_{ebcd}R_{af}R_{ef} + R_{afgh}R_{efgh}R_{ce}\eta_{bd}), \\
\delta B_{11} &= (\delta R_{abcd})(-R_{ebcd}R_{afeg}R_{fg} - \frac{1}{2}R_{aefg}R_{cefg}R_{bd} - \frac{1}{2}R_{eghi}R_{fghi}R_{eafc}\eta_{bd}), \\
\delta B_{12} &= (\delta R_{abcd})(R_{ebcd}R_{ae}R + \frac{1}{2}R_{aefg}R_{cefg}\eta_{bd}R + \frac{1}{2}R_{eghi}R_{fghi}R_{ef}\eta_{ac}\eta_{bd}), \\
\delta B_{13} &= (\delta R_{abcd})(2R_{ebfd}R_{ac}R_{ef} + R_{agch}R_{egfh}R_{ef}\eta_{bd}), \\
\delta B_{14} &= (\delta R_{abcd})(R_{abeg}R_{cdfg}R_{ef} + 2R_{abef}R_{efgd}R_{cg} + R_{efgh}R_{efai}R_{ghci}\eta_{bd}), \\
\delta B_{15} &= (\delta R_{abcd})(R_{aecg}R_{bfdg}R_{ef} + 2R_{aecf}R_{egfd}R_{bg} + R_{efgh}R_{eagi}R_{fchi}\eta_{bd}), \\
\delta B_{16} &= (\delta R_{abcd})(3R_{abef}R_{cdef}R + R_{efgh}R_{efij}R_{ghij}\eta_{ac}\eta_{bd}), \\
\delta B_{17} &= (\delta R_{abcd})(3R_{aecf}R_{bedf}R + R_{efgh}R_{eigj}R_{fihj}\eta_{ac}\eta_{bd}), \\
\delta B_{18} &= (\delta R_{abcd})(R_{ac}R_{bd}R + 2R_{aecf}R_{ef}\eta_{bd}R + R_{efgh}R_{eg}R_{fh}\eta_{ac}\eta_{bd}), \\
\delta B_{19} &= (\delta R_{abcd})(2R_{cdef}R_{ae}R_{bf} + 2R_{aegh}R_{cfgh}R_{ef}\eta_{bd}), \\
\delta B_{20} &= (\delta R_{abcd})(2R_{ebfd}R_{ae}R_{cf} + 2R_{ageh}R_{cgfh}R_{ef}\eta_{bd}), \\
\delta B_{21} &= (\delta R_{abcd})(R_{ae}R_{ce}R_{bd} + 2R_{afeg}R_{ce}R_{fg}\eta_{bd} + R_{ebfd}R_{eg}R_{fg}\eta_{ac}), \\
\delta B_{22} &= 4(\delta R_{abcd})R_{ac}R_{ef}R_{ef}\eta_{bd}, \\
\delta B_{23} &= (\delta R_{abcd})(2R_{ac}\eta_{bd}R^2 + 2R_{ef}R_{ef}\eta_{ac}\eta_{bd}R), \\
\delta B_{24} &= 4(\delta R_{abcd})R_{ef}R_{ae}R_{cf}\eta_{bd}, \\
\delta B_{25} &= (\delta R_{abcd})(3R_{ae}R_{ce}\eta_{bd}R + R_{fg}R_{ef}R_{eg}\eta_{ac}\eta_{bd}), \\
\delta B_{26} &= 4(\delta R_{abcd})\eta_{ac}\eta_{bd}R^3.
\end{aligned} \tag{71}$$

And as like the eq. (12), we define Y tensor as

$$\begin{aligned}
Y_{abcd} &= \frac{1}{2}(Y'_{[ab][cd]} + Y'_{[cd][ab]}), \tag{72} \\
Y'_{abcd} &= b_8(2R_{abcd}R_{ef}R_{ef} + 2R_{efgh}R_{efgh}R_{ac}\eta_{bd}) + b_9(2R_{abcd}R^2 + 2R_{efgh}R_{efgh}\eta_{ac}\eta_{bd}R) \\
&\quad + b_{10}(R_{ebcd}R_{af}R_{ef} + R_{afgh}R_{efgh}R_{ce}\eta_{bd}) \\
&\quad + b_{11}(-R_{ebcd}R_{afeg}R_{fg} - \frac{1}{2}R_{aefg}R_{cefg}R_{bd} - \frac{1}{2}R_{eghi}R_{fghi}R_{eafc}\eta_{bd}) \\
&\quad + b_{12}(R_{ebcd}R_{ae}R + \frac{1}{2}R_{aefg}R_{cefg}\eta_{bd}R + \frac{1}{2}R_{eghi}R_{fghi}R_{ef}\eta_{ac}\eta_{bd}) \\
&\quad + b_{13}(2R_{ebfd}R_{ac}R_{ef} + R_{agch}R_{efgh}R_{ef}\eta_{bd}) \\
&\quad + b_{14}(R_{abeg}R_{cdfg}R_{ef} + 2R_{abef}R_{efgd}R_{cg} + R_{efgh}R_{efai}R_{ghci}\eta_{bd}) \\
&\quad + b_{15}(R_{aecg}R_{bfdg}R_{ef} + 2R_{aecf}R_{egfd}R_{bg} + R_{efgh}R_{eagi}R_{fchi}\eta_{bd}) \\
&\quad + b_{16}(3R_{abef}R_{cdef}R + R_{efgh}R_{efij}R_{ghij}\eta_{ac}\eta_{bd}) \\
&\quad + b_{17}(3R_{aecf}R_{bedf}R + R_{efgh}R_{eigj}R_{fihj}\eta_{ac}\eta_{bd}) \\
&\quad + b_{18}(R_{ac}R_{bd}R + 2R_{aecf}R_{ef}\eta_{bd}R + R_{efgh}R_{eg}R_{fh}\eta_{ac}\eta_{bd}) \\
&\quad + b_{19}(2R_{cdef}R_{ae}R_{bf} + 2R_{aegh}R_{c}R_{fgh}R_{ef}\eta_{bd}) \\
&\quad + b_{20}(2R_{ebfd}R_{ae}R_{cf} + 2R_{ageh}R_{cg}R_{fh}R_{ef}\eta_{bd}) \\
&\quad + b_{21}(R_{ae}R_{ce}R_{bd} + 2R_{afeg}R_{ce}R_{fg}\eta_{bd} + R_{ebfd}R_{eg}R_{fg}\eta_{ac}) \\
&\quad + 4b_{22}R_{ac}R_{ef}R_{ef}\eta_{bd} + b_{23}(2R_{ac}\eta_{bd}R^2 + 2R_{ef}R_{ef}\eta_{ac}\eta_{bd}R) \\
&\quad + 4b_{24}R_{ef}R_{ae}R_{cf}\eta_{bd} + b_{25}(3R_{ae}R_{ce}\eta_{bd}R + R_{fg}R_{ef}R_{eg}\eta_{ac}\eta_{bd}) + 4b_{26}\eta_{ac}\eta_{bd}R^3.
\end{aligned}$$

Following the similar calculations in the eq. (11), finally we obtain generic equations of motion

$$\begin{aligned}
E_{ij} &\equiv R_{ij} - \frac{1}{2}\eta_{ij}R + \gamma\ell_s^{12} \left\{ -\frac{1}{2}\eta_{ij} \left(t_8 t_8 R^4 - \frac{1}{4!} \epsilon_{11} \epsilon_{11} R^4 + \sum_{n=8}^{26} b_n B_n \right) \right. \\
&\quad + \frac{3}{2}R_{abci}X^{abc}{}_j - \frac{1}{2}R_{abcj}X^{abc}{}_i - 2D_{(a}D_{b)}X^a{}_{ij}{}^b \\
&\quad \left. + \frac{3}{2}R_{abci}Y^{abc}{}_j - \frac{1}{2}R_{abcj}Y^{abc}{}_i - 2D_{(a}D_{b)}Y^a{}_{ij}{}^b \right\} = 0. \tag{73}
\end{aligned}$$

In order to evaluate these equations, we need to insert the values of the Riemann tensor (62) into the above. Since the Ricci tensor and the scalar curvature become zero we obtain $B_n = 0$, and parts of b_{11} , b_{14} , b_{15} , b_{16} and b_{17} in the Y tensor only contribute to the above equations of motion.

Below we repeat the similar calculations in the appendix B. Each component of Y_{abcd} is

evaluated as

$$\begin{aligned}
Y_{0101} &= \frac{1}{U_0^6 x^{27} \ell_s^{12}} \left\{ \frac{11907}{2} b_{11} (1 + x^7) - 21609 b_{14} (1 + x^7) \right. \\
&\quad \left. - \frac{3087}{2} b_{15} (1 + x^7) - 85176 b_{16} - 10458 b_{17} \right\}, \\
Y_{0\bar{a}0\bar{a}} &= \frac{1}{U_0^6 x^{27} \ell_s^{12}} \left\{ \frac{11907}{8} (-1 + 4x^7) b_{11} + \frac{63}{4} (5 - 1372x^7) b_{14} \right. \\
&\quad \left. - \frac{63}{4} (17 + 98x^7) b_{15} - 85176 b_{16} - 10458 b_{17} \right\}, \\
Y_{011\ddagger} &= \frac{\sqrt{x^7 - 1}}{U_0^6 x^{\frac{47}{2}} \ell_s^{12}} \left\{ -\frac{11907}{2} b_{11} + 21609 b_{14} + \frac{3087}{2} b_{15} \right\}, \\
Y_{0\bar{a}\bar{a}\ddagger} &= \frac{\sqrt{x^7 - 1}}{U_0^6 x^{\frac{47}{2}} \ell_s^{12}} \left\{ -\frac{11907}{2} b_{11} + 21609 b_{14} + \frac{3087}{2} b_{15} \right\}, \\
Y_{0\ddagger0\ddagger} &= \frac{1}{U_0^6 x^{27} \ell_s^{12}} \left\{ \frac{11907}{2} b_{11} - 21609 b_{14} - \frac{3087}{2} b_{15} - 85176 b_{16} - 10458 b_{17} \right\} \quad (74) \\
Y_{1\ddagger1\ddagger} &= \frac{1}{U_0^6 x^{27} \ell_s^{12}} \left\{ -\frac{11907}{2} b_{11} (2 - x^7) + 21609 (2 - x^7) b_{14} \right. \\
&\quad \left. + \frac{3087}{2} b_{15} (2 - x^7) + 85176 b_{16} + 10458 b_{17} \right\}, \\
Y_{1\bar{a}1\bar{a}} &= \frac{1}{U_0^6 x^{27} \ell_s^{12}} \left\{ -\frac{35721}{8} b_{11} + \frac{86121}{4} b_{14} + \frac{7245}{4} b_{15} + 85176 b_{16} + 10458 b_{17} \right\}, \\
Y_{\bar{a}\ddagger\bar{a}\ddagger} &= \frac{1}{U_0^6 x^{27} \ell_s^{12}} \left\{ \frac{11907}{8} b_{11} (-3 + 4x^7) + \frac{63}{4} b_{14} (1367 - 1372x^7) \right. \\
&\quad \left. + \frac{63}{4} b_{15} (115 - 98x^7) + 85176 b_{16} + 10458 b_{17} \right\}, \\
Y_{\bar{a}\bar{b}\bar{a}\bar{b}} &= \frac{1}{U_0^6 x^{27} \ell_s^{12}} \left\{ \frac{11907}{4} b_{11} - \frac{315}{2} b_{14} + \frac{1071}{2} b_{15} + 85176 b_{16} + 10458 b_{17} \right\},
\end{aligned}$$

where $\bar{a}, \bar{b} = 2, \dots, 9$. By using these results it is possible to evaluate the higher derivative terms which depend on the Y tensor in the eq. (73). The RY terms are calculated as

$$\begin{aligned}
R_{abc0} Y^{abc}_0 &= \frac{1}{U_0^8 x^{29} \ell_s^{16}} \left\{ 416745 b_{11} - 1214514 b_{14} - 71442 b_{15} \right\}, \\
R_{abc1} Y^{abc}_1 &= \frac{1}{U_0^8 x^{36} \ell_s^{16}} \left\{ -416745 b_{11} + 1214514 b_{14} + 71442 b_{15} \right\}, \\
R_{abc\ddagger} Y^{abc}_{\ddagger} &= \frac{x^7 - 1}{U_0^8 x^{36} \ell_s^{16}} \left\{ 416745 b_{11} - 1214514 b_{14} - 71442 b_{15} \right\}, \quad (75) \\
R_{abc\bar{a}} Y^{abc}_{\bar{a}} &= \frac{1}{4U_0^8 x^{36} \ell_s^{16}} \delta_{\bar{a}\bar{b}} \left\{ 416745 b_{11} - 1214514 b_{14} - 71442 b_{15} \right\}, \\
R_{abc0} Y^{abc}_{\ddagger} &= R_{abc\ddagger} Y^{abc}_0 = \frac{\sqrt{x^7 - 1}}{U_0^8 x^{\frac{65}{2}} \ell_s^{16}} \left\{ 416745 b_{11} - 1214514 b_{14} - 71442 b_{15} \right\},
\end{aligned}$$

and DDY terms become

$$\begin{aligned}
D_{(a}D_b)Y^a{}_{00}{}^b &= \frac{1701}{U_0^8 x^{36} \ell_s^{16}} \left\{ -\frac{7}{2}(-459 - 235x^7 + 540x^{14})b_{11} \right. \\
&\quad + (-6507 - 2397x^7 + 6860x^{14})b_{14} + \frac{1}{2}(-999 - 282x^7 + 980x^{14})b_{15} \\
&\quad \left. + (-36504 + 31772x^7)b_{16} + (-4482 + 3901x^7)b_{17} \right\}, \\
D_{(a}D_b)Y^a{}_{11}{}^b &= \frac{1701}{U_0^8 x^{36} \ell_s^{16}} \left\{ -7(31 + 46x^7)b_{11} + 4(6 + 505x^7)b_{14} \right. \\
&\quad \left. + \frac{1}{2}(-75 + 376x^7)b_{15} + 676(-9 + 16x^7)b_{16} + 83(-9 + 16x^7)b_{17} \right\}, \\
D_{(a}D_b)Y^a{}_{\ddot{1}\ddot{1}}{}^b &= \frac{1701}{U_0^8 x^{36} \ell_s^{16}} \left\{ -\frac{7}{2}(1034 - 1455x^7 + 540x^{14})b_{11} \right. \\
&\quad + (13724 - 18897x^7 + 6860x^{14})b_{14} + \frac{1}{2}(2021 - 2742x^7 + 980x^{14})b_{15} \\
&\quad \left. + 676(47 - 33x^7)b_{16} + 83(47 - 33x^7)b_{17} \right\}, \tag{76} \\
D_{(a}D_b)Y^a{}_{\bar{a}\bar{b}}{}^b &= \frac{-4 + 3x^7}{U_0^8 x^{36} \ell_s^{16}} \delta_{\bar{a}\bar{b}} \left\{ \frac{1917027}{4}b_{11} - \frac{6013035}{2}b_{14} - \frac{559629}{2}b_{15} \right. \\
&\quad \left. - 16098264b_{16} - 1976562b_{17} \right\}, \\
D_{(a}D_b)Y^a{}_{0\ddot{1}}{}^b &= D_{(a}D_b)Y^a{}_{\ddot{1}0}{}^b \\
&= \frac{\sqrt{x^7 - 1}}{U_0^8 x^{\frac{65}{2}} \ell_s^{16}} \left\{ \frac{59535}{2}(115 - 108x^7)b_{11} - 11907(1031 - 980x^7)b_{14} \right. \\
&\quad \left. - 11907(73 - 70x^7)b_{15} + 8049132b_{16} + 988281b_{17} \right\}.
\end{aligned}$$

As mentioned before, only b_{11} , b_{14} , b_{15} , b_{16} and b_{17} appeared in the calculations.

By using the ansatz (14) and inserting values of X and Y tensors into the equations of

motion (73), we obtain five independent equations with parameters b_{11} , b_{14} , b_{15} , b_{16} and b_{17} .

$$\begin{aligned}
E_1 = & -63x^{34}f_1 - 9x^{35}f_1' - 49x^{41}h_1 + 49x^{34}(1-x^7)h_2 + 23x^{35}(1-x^7)h_2' + 2x^{36}(1-x^7)h_2'' \\
& + 98x^{41}h_3 + 7x^{42}h_3' - 63402393600x^{14} + 70230343680x^7 + 1062512640 \\
& + (25719120b_{11} - 93350880b_{14} - 6667920b_{15})x^{14} \\
& + (-9525600b_{11} + 27760320b_{14} + 1632960b_{15} - 432353376b_{16} - 53084808b_{17})x^7 \\
& - 21861252b_{11} + 88547256b_{14} + 6797196b_{15} + 496746432b_{16} + 60991056b_{17} = 0,
\end{aligned} \tag{77}$$

$$\begin{aligned}
E_2 = & 63x^{34}f_1 + 9x^{35}f_1' + 7x^{34}(9-2x^7)h_1 + 9x^{35}(1-x^7)h_1' - 112x^{34}(1-x^7)h_2 \\
& - 16x^{35}(1-x^7)h_2' - 98x^{41}h_3 - 7x^{42}h_3' - 2159861760x^7 - 5730600960 \\
& + (4381776b_{11} - 27488160b_{14} - 2558304b_{15} - 147184128b_{16} - 18071424b_{17})x^7 \\
& + 1285956b_{11} + 4531464b_{14} + 796068b_{15} + 82791072b_{16} + 10165176b_{17} = 0,
\end{aligned} \tag{78}$$

$$\begin{aligned}
E_3 = & 133x^{34}f_1 + 35x^{35}f_1' + 2x^{36}f_1'' + 28x^{34}(3-10x^7)h_1 + 7x^{35}(4-7x^7)h_1' + 2x^{36}(1-x^7)h_1'' \\
& - 7x^{34}(5-26x^7)h_2 - 21x^{35}(1-2x^7)h_2' - 2x^{36}(1-x^7)h_2'' + 98x^{41}h_3 + 7x^{42}h_3' \\
& + 5669637120x^7 - 8626383360 \\
& + (-11502162b_{11} + 72156420b_{14} + 6715548b_{15} + 386358336b_{16} + 47437488b_{17})x^7 \\
& + 15752961b_{11} - 97423074b_{14} - 9025506b_{15} - 515144448b_{16} - 63249984b_{17} = 0,
\end{aligned} \tag{79}$$

$$\begin{aligned}
E_4 = & 259x^{34}f_1 + 53x^{35}f_1' + 2x^{36}f_1'' + 147x^{34}(1-3x^7)h_1 + x^{35}(37-58x^7)h_1' \\
& + 2x^{36}(1-x^7)h_1'' + 147x^{41}h_2 + 21x^{42}h_2' + 294x^{41}h_3 + 21x^{42}h_3' \\
& - 63402393600x^{14} + 133632737280x^7 - 71292856320 \\
& + x^{14}(25719120b_{11} - 93350880b_{14} - 6667920b_{15}) \\
& + x^7(-67631760b_{11} + 252292320b_{14} + 18370800b_{15} + 303567264b_{16} + 37272312b_{17}) \\
& + 47580372b_{11} - 181898136b_{14} - 13465116b_{15} - 432353376b_{16} - 53084808b_{17} = 0,
\end{aligned} \tag{80}$$

$$\begin{aligned}
E_5 = & 49x^{34}h_1 + 7x^{35}h_1' + 49x^{34}h_2 - x^{35}h_2' - x^{36}h_2'' - 98x^{34}h_3 - 22x^{35}h_3' - x^{36}h_3'' \\
& - 63402393600x^7 + 70230343680 \\
& + x^7(25719120b_{11} - 93350880b_{14} - 6667920b_{15}) \\
& - 25719120b_{11} + 93350880b_{14} + 6667920b_{15} - 64393056b_{16} - 7906248b_{17} = 0.
\end{aligned} \tag{81}$$

Here we defined $E_1 = 4U_0^8 \ell_s^4 x^{36} \gamma^{-1} E_{00}$, $E_2 = 4U_0^8 \ell_s^4 x^{36} \gamma^{-1} E_{11}$, $E_3 = 4U_0^8 \ell_s^4 x^{36} \gamma^{-1} E_{22}$, $E_4 = 4U_0^8 \ell_s^4 x^{36} \gamma^{-1} E_{\ddagger\ddagger}$ and $E_5 = 4U_0^8 \ell_s^4 x^{\frac{65}{2}} (-1+x^7)^{-\frac{1}{2}} \gamma^{-1} E_{0\ddagger}$.

The equations (77)–(81) can be solved by following the details in the section 4. And the

final form of the solution becomes

$$\begin{aligned}
h_1 &= \left(-\frac{440559}{4}b_{11} + \frac{768775}{2}b_{14} + \frac{53333}{2}b_{15} + 927472b_{16} + 113876b_{17} + \frac{1302501760}{9} \right) \frac{1}{x^{34}} \\
&\quad + (23814b_{11} - 86436b_{14} - 6174b_{15} - 57462496) \frac{1}{x^{27}} + \frac{12051648}{13x^{20}} - \frac{4782400}{13x^{13}} \\
&\quad - \frac{3747840}{x^7} + \frac{4099200}{x^6} - \frac{1639680(x-1)}{(x^7-1)} + 117120 \left(18 - \frac{23}{x^7} \right) I(x), \\
h_2 &= \left(-\frac{11907}{4}b_{11} + \frac{315}{2}b_{14} - \frac{1071}{2}b_{15} - 170352b_{16} - 20916b_{17} + 19160960 \right) \frac{1}{x^{34}} \\
&\quad + (23814b_{11} - 86436b_{14} - 6174b_{15} - 58528288) \frac{1}{x^{27}} + \frac{2213568}{13x^{20}} - \frac{1229760}{13x^{13}} \\
&\quad - \frac{2108160}{x^7} + \frac{2459520}{x^6} + 1054080 \left(2 - \frac{1}{x^7} \right) I(x), \tag{82} \\
h_3 &= \left(-\frac{11907}{4}b_{11} + \frac{76027}{2}b_{14} + \frac{8225}{2}b_{15} - 94640b_{16} - 11620b_{17} + \frac{361110400}{9} \right) \frac{1}{x^{34}} \\
&\quad + (23814b_{11} - 86436b_{14} - 6174b_{15} - 59840032) \frac{1}{x^{27}} - \frac{24021312}{13x^{20}} \\
&\quad - \frac{58072000}{13x^{13}} - \frac{2108160}{x^7} + \frac{2459520}{x^6} + 117120 \left(18 - \frac{41}{x^7} \right) I(x), \\
f_1 &= \left(\frac{440559}{4}b_{11} - \frac{730919}{2}b_{14} - \frac{48685}{2}b_{15} - 889616b_{16} - 109228b_{17} - \frac{1208170880}{9} \right) \frac{1}{x^{34}} \\
&\quad + \left(-130977b_{11} + 432810b_{14} + 28728b_{15} + 1022112b_{16} + 125496b_{17} + 161405664 \right) \frac{1}{x^{27}} \\
&\quad + \frac{5738880}{13x^{20}} + \frac{956480}{x^{13}} + \frac{819840}{x^7} I(x).
\end{aligned}$$

The function $I(x)$ is given by the eq. (34) and integral constants are determined so as to satisfy that $h_i(1)$ are finite and $h_i(x), f_1(x) \sim \mathcal{O}(x^{-8})$ when x goes to the infinity. Notice that $b_{11}, b_{14}, b_{15}, b_{16}$ and b_{17} only appear in the coefficients of x^{-27} and x^{-34} . The solution is reliable up to $\mathcal{O}(\epsilon^2)$.

D Thermodynamics of Black 0-Brane with Generic R^4 Terms

In this appendix, we examine thermodynamics of the quantum near horizon geometry of the black 0-brane (82) by following the arguments in the section 5. Although the solution is modified, the results obtained until the eq. (50) do not change. Since the effective action is modified as in the eq. (70), the eq. (51) should be replaced with

$$\frac{\partial S_{11}}{\partial R_{\mu\nu\rho\sigma}} = \frac{1}{2\kappa_{11}^2} \{ g^{\mu[\rho} g^{\sigma]\nu} + \gamma \ell_s^{12} (X^{\mu\nu\rho\sigma} + Y^{\mu\nu\rho\sigma}) \}. \tag{83}$$

The entropy of the quantum near horizon geometry of the black 0-brane is evaluated as

$$\begin{aligned}
S &= \frac{4\pi}{2\kappa_{11}^2} \int_{\text{H}} d\Omega_8 dz \sqrt{h} \left(1 - \frac{1}{2} \gamma \ell_s^{12} (X^{\mu\nu\rho\sigma} + Y^{\mu\nu\rho\sigma}) N_{\mu\nu} N_{\rho\sigma} \right) \\
&= \frac{4\pi}{2\kappa_{11}^2} \int_{\text{H}} d\Omega_8 dz \sqrt{h} \left(1 - 2\gamma \ell_s^{20} U_0^2 H_1^{-1} (X^{txtx} + Y^{txtx}) \right) \\
&= \frac{4\pi}{2\kappa_{11}^2} \int_{\text{H}} d\Omega_8 dz \sqrt{h} \left\{ 1 + \epsilon U_0^{-6} (40642560 \right. \\
&\quad \left. - 23814b_{11} + 86436b_{14} + 6174b_{15} + 170352b_{16} + 20916b_{17}) \right\} \\
&= \frac{4}{49} a_1 N^2 \tilde{U}_0^{\frac{9}{2}} \left\{ 1 + \epsilon \left(-\frac{9}{14} f_1(1) + \frac{1}{2} h_2(1) + 40642560 \right. \right. \\
&\quad \left. \left. - 23814b_{11} + 86436b_{14} + 6174b_{15} + 170352b_{16} + 20916b_{17} \right) \tilde{U}_0^{-6} \right\} \\
&= \frac{4}{49} a_1^{-\frac{4}{5}} N^2 \tilde{T}^{\frac{9}{5}} \left\{ 1 + \epsilon a_1^{\frac{12}{5}} \left(-\frac{9}{5} f_1(1) - \frac{9}{35} f_1'(1) + \frac{9}{10} h_1(1) + \frac{1}{2} h_2(1) + 40642560 \right. \right. \\
&\quad \left. \left. - 23814b_{11} + 86436b_{14} + 6174b_{15} + 170352b_{16} + 20916b_{17} \right) \tilde{T}^{-\frac{12}{5}} \right\} \\
&= a_3 N^2 \tilde{T}^{\frac{9}{5}} (1 + \epsilon a_5 \tilde{T}^{-\frac{12}{5}}). \tag{84}
\end{aligned}$$

Notice that $f_1(1)$, $f_1'(1)$, $h_1(1)$ and $h_2(1)$ depend on b_{11} , b_{14} , b_{15} , b_{16} and b_{17} . The value of a_3 is given in the section 5, and a_5 is given by

$$\begin{aligned}
a_5 &= a_1^{\frac{12}{5}} \left(-\frac{9}{5} f_1(1) - \frac{9}{35} f_1'(1) + \frac{9}{10} h_1(1) + \frac{1}{2} h_2(1) + 40642560 \right. \\
&\quad \left. - 23814b_{11} + 86436b_{14} + 6174b_{15} + 170352b_{16} + 20916b_{17} \right). \tag{85}
\end{aligned}$$

It seems that a_5 depends on b_{11} , b_{14} , b_{15} , b_{16} and b_{17} . The explicit calculation, however, shows that $a_5 = a_4$ and the result does not depend on the ambiguities of the effective action. Thus the physical quantities of the black 0-brane are free from the ambiguities and uniquely determined.

References

- [1] J. Polchinski, “*Dirichlet branes and Ramond-Ramond charges*”, Phys. Rev. Lett. **75** (1995) 4724.
- [2] G. W. Gibbons and K. Maeda, “*Black holes and membranes in higher dimensional theories with dilaton fields*”, Nucl. Phys. **B298** (1988) 741.
- [3] G. T. Horowitz and A. Strominger, “*Black strings and p-branes*”, Nucl. Phys. **B360** (1991) 197.
- [4] A. Strominger and C. Vafa, “*Microscopic origin of the Bekenstein-Hawking entropy*”, Phys. Lett. **B379** (1996) 99.

- [5] J. M. Maldacena, “*The Large N limit of superconformal field theories and supergravity*”, Adv. Theor. Math. Phys. **2** (1998) 231.
- [6] S. Gubser, I. R. Klebanov, and A. M. Polyakov, “*Gauge theory correlators from non-critical string theory*”, Phys. Lett. **B428** (1998) 105.
- [7] E. Witten, “*Anti-de Sitter space and holography*”, Adv. Theor. Math. Phys. **2** (1998) 253.
- [8] N. Itzhaki, J. M. Maldacena, J. Sonnenschein and S. Yankielowicz, “*Supergravity and the large N limit of theories with sixteen supercharges*”, Phys. Rev. **D58** (1998) 046004.
- [9] E. Witten, “*Bound states of strings and p -branes*”, Nucl. Phys. **B460** (1996) 335.
- [10] B. de Wit, J. Hoppe, and H. Nicolai, “*On quantum mechanics of supermembranes*”, Nucl. Phys. **B305** (1988) 545.
- [11] T. Banks, W. Fischler, S. H. Shenker and L. Susskind, “ *M theory as a matrix model: A conjecture*”, Phys. Rev. **D55** (1997) 5112.
- [12] P. K. Townsend, “*The eleven-dimensional supermembrane revisited*”, Phys. Lett. **B350** (1995) 184.
- [13] E. Witten, “*String theory dynamics in various dimensions*”, Nucl. Phys. **B443** (1995) 85.
- [14] D. Kabat, G. Lifschytz and D. A. Lowe, “*Black Hole Thermodynamics from Calculations in Strongly Coupled Gauge Theory*”, Phys. Rev. Lett. **86** (2001) 1426.
- [15] M. Hanada, J. Nishimura and S. Takeuchi, “*Non-lattice simulation for supersymmetric gauge theories in one dimension*”, Phys. Rev. Lett. **99** (2007) 161602.
- [16] S. Catterall and T. Wiseman, “*Towards lattice simulation of the gauge theory duals to black holes and hot strings*”, JHEP **12** (2007) 104.
- [17] S. Catterall and T. Wiseman, “*Black hole thermodynamics from simulations of lattice Yang-Mills theory*”, Phys. Rev. **D78** (2008) 041502.
- [18] M. Hanada, A. Miwa, J. Nishimura and S. Takeuchi, “*Schwarzschild radius from Monte Carlo calculation of the Wilson loop in supersymmetric matrix quantum mechanics*”, Phys. Rev. Lett. **102** (2009) 181602.

- [19] M. Hanada, Y. Hyakutake, J. Nishimura and S. Takeuchi, “*Higher derivative corrections to black hole thermodynamics from supersymmetric matrix quantum mechanics*”, Phys. Rev. Lett. **102** (2009) 191602.
- [20] M. Hanada, J. Nishimura, Y. Sekino and T. Yoneya, “*Monte Carlo studies of Matrix theory correlation functions*”, Phys. Rev. Lett. **104** (2010) 151601.
- [21] S. Catterall and T. Wiseman, “*Extracting black hole physics from the lattice*”, JHEP **1004** (2010) 077.
- [22] M. Hanada, J. Nishimura, Y. Sekino and T. Yoneya, “*Direct test of the gauge-gravity correspondence for Matrix theory correlation functions*”, JHEP **1112** (2011) 020.
- [23] D. Kadoh and S. Kamata, “*One dimensional supersymmetric Yang-Mills theory with 16 supercharges*”, PoS LATTICE **2012** (2012) 064.
- [24] M. Hanada, Y. Hyakutake, G. Ishiki and J. Nishimura, “*Holographic description of quantum black hole on a computer*”, arXiv:1311.5607.
- [25] S. Catterall, D. B. Kaplan and M. Unsal, “*Exact lattice supersymmetry*”, Phys. Rept. **484** (2009) 71.
- [26] J. Nishimura, “*The origin of space-time as seen from matrix model simulations*”, Prog. Theor. Exp. Phys. **2012** (2012) 01A101.
- [27] D. J. Gross and E. Witten, “*Superstring Modifications of Einstein’s Equations*”, Nucl. Phys. **B277** (1986) 1.
- [28] M. de Roo, H. Suelmann and A. Wiedemann, “*Supersymmetric R^4 -actions in ten dimensions*” Phys. Lett. **B280** (1992) 39.
- [29] M. de Roo, H. Suelmann and A. Wiedemann, “*The Supersymmetric effective action of the heterotic string in ten-dimensions*” Nucl. Phys. **B405** (1993) 326.
- [30] H. Suelmann, “*String effective actions and supersymmetry*”, PhD Thesis, Groningen University, 1994.
- [31] A. A. Tseytlin, “ *R^4 -terms in 11-dimensions and Conformal anomaly of $(2,0)$ theory*”, Nucl. Phys. **B584** (2000) 233.
- [32] K. Becker and M. Becker, “*Supersymmetry breaking, M theory and fluxes*”, JHEP **0107** (2001) 038.

- [33] K. Peeters, P. Vanhove and A. Westerberg, “*Supersymmetric higher-derivative actions in ten and eleven dimensions, the associated superalgebras and their formulation in superspace*” *Class. Quant. Grav.* **18** (2001) 843.
- [34] G. Policastro and D. Tsimpis, “ *R^4 , purified*”, *Class. Quant. Grav.* **23** (2006) 4753.
- [35] Y. Hyakutake and S. Ogushi, “Higher derivative corrections to eleven dimensional supergravity via local supersymmetry”, *JHEP* **0602** (2006) 068.
- [36] Y. Hyakutake, “Toward the Determination of R^3F^2 Terms in M-theory”, *Prog. Theor. Phys.* **118** (2007) 109.
- [37] A. Dabholkar, “Exact counting of black hole microstates”, *Phys. Rev. Lett.* **94** (2005) 241301.
- [38] H. Ooguri, A. Strominger, C. Vafa, “Black hole attractors and the topological string”, *Phys. Rev.* **D70** (2004) 106007.
- [39] E. Cremmer, B. Julia and J. Scherk, “Supergravity Theory in Eleven-Dimensions”, *Phys. Lett.* **B76** (1978) 409.
- [40] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, “Large N field theories, string theory and gravity”, *Phys. Rept.* **323** (2000) 183.
- [41] R. M. Wald, “*Black Hole Entropy is Noether Charge*”, *Phys. Rev.* **D48** (1993) R3427.
- [42] V. Iyer and R.M. Wald, “*Some Properties of Noether Charge and a Proposal for Dynamical Black Hole Entropy*”, *Phys. Rev.* **D50** (1994) 846.
- [43] Y.-H. Lin, S.-H. Shao, Y. Wang and X. Yin, “*A Low Temperature Expansion for Matrix Quantum Mechanics*”, arXiv:1304.1593.
- [44] T. Morita, S. Shiba, T. Wiseman and B. Withers, “*Warm p-soup and near extremal black holes*”, arXiv:1311.6540.
- [45] S. L. Braunstein, S. Pirandola and K. Zyczkowski, “*Better Late than Never: Information Retrieval from Black Holes*”, *Phys. Rev. Lett.* **110** (2013) 101301.
- [46] A. Almheiri, D. Marolf, J. Polchinski and J. Sully, “*Black holes: Complementarity or firewalls?*”, *JHEP* **1302** (2013) 062.